



International Journal of Medical and All Body Health Research

Antibacterial activity of *Acacia nilotica* and *Jatropha curcas* plants leave extracts against multi-drug resistant strains of *Pseudomonas aeruginosa* and *Staphylococcus aureus*

Hassan AU ^{1*}, Garba S ², MS Buhari ³, Appah J ⁴, UO Ozojiofor ⁵, Egbe NE ⁶

^{1, 3, 5, 6} Department of Biotechnology, Nigerian Defence Academy, Kaduna, Nigeria

² Department of Chemistry, Nigerian Defence Academy, Kaduna, Nigeria

⁴ Department of Biology, Nigerian Defence Academy, Kaduna, Nigeria

* Corresponding Author: Hassan AU

Article Info

ISSN (online): 2582-8940

Volume: 04

Issue: 03

July-September 2023

Received: 10-08-2023;

Accepted: 08-09-2023

Page No: 34-46

Abstract

With an elevated increase in the emergence of multidrug resistance strains of different kinds of virulent Pathogenic Bacteria to various kinds of antibiotics worldwide, which has ultimately become a serious medical and economic public health issues in Nigeria and other African Countries. It however remains a serious medical, economic and social problem. This necessitates the need for finding novel antimicrobial agents, especially from natural systems like ethno-medicinal plants with minimal pathogenic resistance and therapeutic side effects. This research is aimed at evaluating the antibacterial activity of *Acacia nilotica* and *Jatropha curcas* ethno-medicinal plants against multi-drug resistant strains of *Pseudomonas aeruginosa* and *Staphylococcus aureus*. *A. nilotica* and *J. curcas* leaves were extracted using chloroform, to get the crude extracts of the various plants material. One hundred and fifty (150) clinical and environmental samples each, were collected for the isolation of *Pseudomonas aeruginosa* and *Staphylococcus aureus* each. The antibiogram profile and resistance level of the isolated strains was ascertain using the standard disc diffusion bioassay. Multi-drug resistance genes; BLA-TEM, BLA-SHV and CTX-M presence were evaluated and also served as another form of selection protocol for extremely resistant strains of the selected pathogenic bacterial species. In-vitro and in-vivo bioassay was done to evaluate the antimicrobial potential of the plants extracts on the most resistant strains of the two pathogenic bacteria. Tannins, saponin, alkaloids, flavonoids, phenols, terpenoids, quinones, phytosterols and phlobatanins were found present in these plants extracts. Eleven *Pseudomonas aeruginosa* and four *Staphylococcus aureus* extreme multi-drug resistance strains were successfully isolated and identified. The in-vitro bioassay reveals that *J. curcas* extracts has the highest antimicrobial activity against the various multi-drug resistant strains, while *A. nilotica* extract also maintains some level of activity against the resistant strains. The in-vivo bioassay reconfirmed the antibacterial effects of the plants, against all the resistant bacteria. Decoction of the plants extracts also proves an even more effective activity on the resistant strains of both bacteria. *J. curcas* and *A. nilotica* plant extracts could provide an alternative pharmaceuticals against multidrug resistant strains of *Pseudomonas aeruginosa* and *Staphylococcus aureus*.

Keywords: Resistance, bioassay, antimicrobial, ethno-medicinal plant, phytometabolites, bacteria

Introduction

Floral organisms are potential source of medical agents, as they are traditionally used in treatment and management many diseases and infections of different categories ^[1]. Nowadays, the herbal medicines are still widely used in conventional and alternative medical practices in developed and developing countries as a complementary medicine. Plants are been used as a source of inspiration in the development of novel drug or part of it ^[2].

Phytochemicals are metabolic chemical compounds that occur naturally in plants, which are responsible for a lot of plants peculiar features and capability, such as; color and organoleptic properties, like the deep purple of blueberries or the smell of garlic [3]. Phytochemicals are wide variety of chemical compounds, which can be sorted by their differential chemical class, biosynthetic origin and functional groups [4]. Plant however also used in the treatment and management of zoonotic hazards such as bites from snakes, bees, scorpions and other zoonotic animals [5].

Pseudomonas aeruginosa and *Staphylococcus aureus* are virulent species of bacteria, which are two of the most causative agents of nosocomial infections (hospital acquired infections). They are aerobic, opportunistic pathogens, whose infections are usually very difficult to treat, because of their elevated intrinsic resistance as well as their capacity to acquire resistance to different antibiotics [6]. Bacterial multi-drug resistance genes like BLA-TEM, BLA-SHV and CTX-M genes usually encode production of Extended Spectrum Beta Lactamase, which confers virulent resistance to various forms of Beta-lactams antibiotics among others, this is one of the most important form of antibiotic drug resistance. The Bla-SHV gene together with other functional agents aids in maintaining bacterial genomic integrity, by inducing DNA damage repair mechanism and withstanding environmental stress by the host immune response systems [7]. The extended spectrum beta Lactamase enzymes of the TEM and SHV bacterial producing families are the mutant forms of beta-lactamases, while the CTX-M family initially originated from certain environmental bacteria [8]. Medicinal plants are the biggest bio-resource of drugs of traditional systems of medicine, modern medicines, nutraceuticals, food supplements, folk medicines, pharmaceutical intermediates/semi-antibiotics and chemical entities for synthetic drugs [9]. Medicinal plants like *Jatropha curcas* which is a plant that belongs to the Euphorbiaceae family, which is a perennial, monoecious shrub or small tree up to 6 m high. *Jatropha curcas* is used traditionally in the treatment of wounds and malaria [10]. The plant is also used in controlling soil degradation, erosion, desertification and in increasing soil fertility. Researchers have also reported the use of *Jatropha curcas* oil for production of biodiesel [11]. *Acacia nilotica* is another evergreen plant, with an average sizes of about 2.5 - 25 m, which usually has a short, thick and cylindrical trunk [12]. *Acacia nilotica* (bagaruwa in Hausa) has been designated and used as medicinal plants in parts of Northern Nigeria, West Africa, North Africa and other parts of the world [13]. It's used to treat infections such as diarrhea, dysentery, leprosy, cancers, ulcer and diabetes [14].

Materials and Method

Sample Collection and Processing

Plant Sampling and Authentication

Leaves of *Jatropha curcas* and *Acacia nilotica*, were collected from different parts of Katsina and Kaduna states, Nigeria. The plants materials were then identified and authenticated, properly washed using tap water and rinsed with distilled water, before air drying under shade. Each dried plant sample material was then pulverized separately into powder and stored in clean polythene bags at ambient temperature for further analysis [15].

Isolation of *Pseudomonas aeruginosa*

Pseudomonas aeruginosa isolates were collected from

different hospitals and their surroundings, refuse dumps, mechanic workshops and poultry houses in Kaduna metropolis. The isolates were then further sub-cultured using cetrimide agar, blood agar and mannitol salt agar, where microbiological and biochemically reconfirmations analyses were carried out on the individual isolates, before storage on slants of nutrient agar and refrigerated at 4°C, until required for use [6].

Plant Material Extraction

A portion (500g) each of the respective processed stored powder of the plants materials, were separately percolated in a maceration bottle, using 300 ml of chloroform. The powdered plant sample is allowed to macerate for a period of one week in the solvent inside the percolator. Each extract are then filtered using a pressure suction pump and then evaporated to dryness at 40 °C using rotary evaporator. Individual residue produced are then allowed to cool, weighed and stored in a refrigerator, until use [16].

Phytochemical Screening

Qualitative and quantitative phytochemical analysis were done, using standard protocols, as described by [17].

Microbiological Identification of *Pseudomonas aeruginosa*

Test isolates was inoculated and sub-cultured on different kinds of culture mediums, which includes; nutrient agar, blood agar and the cetrimide agar, for Colony-morphology identification. In addition to macroscopic characteristics, gram-staining and microscopic examination was carried out with special regard towards cell shape and arrangement [18].

Biochemical Identification of *Pseudomonas aeruginosa* isolates

Biochemical analyses was carried out on each of the isolates, including the Nitrate Reduction, Catalase, Oxidase, Methyl-Red, Motility, Indole, Citrate, Voges Proskauer and Urease testes [19].

Susceptibility Test

The antibiotic susceptibility was tested on Mueller–Hinton agar plates, using the standard disk diffusion technique (Kirby-Bauer test). Briefly, the confirmed *P. aeruginosa* are grown on Mueller-Hilton agar and incubated at 37°C for 18–24h. After incubation, few colonies was reconstituted on sterile physiological saline to a diluent approximating to 0.5 McFarland standards. The bacterial suspension was spread onto a Mueller–Hinton agar plate surface to form a confluent lawn and incubated for approximately 15 min, afterwards, the agar plates was impregnated with the following antibiotic discs; Amikacin (30µg), Gentamicin (10µg), Tobramycin (10µg), Ceftazidime (30µg), Cefepime (30µg), Doripenem (30µg), Imipenem (30µg), Meropenem (30µg), Piperacillin (100µg), Ciprofloxacin (5µg), Levofloxacin (5µg), Ofloxacin (5µg) and Piperacillin-tazobactam (100/10µg). Plates were read after 24h of incubation and results was recorded [20].

Preparation of Fraction Stock Solution

5 g of each extract fraction was diluted with 10 mL of 10% dimethyl sulfoxide (DMSO), which serves as stock Solution (500mg/ml), and then sterilized by filtration through a bacterial filter of pore size 0.45µm using positive pressure. Then, filtrate were then kept at 4°C in refrigerator until use [21].

Inoculum Standardization

A looful of the confirmed test isolates was picked using a sterile wire loop and emulsified in 3-4mls of normal saline and properly shaken. The suspension turbidity was then matched with the 0.5 McFarland standard [22].

Identification of Resistance Genes

The evaluation for the presence of resistance genes in the identified isolates of *P. aeruginosa* and *S. aureus* was done using multiplex polymerase chain reaction technique as described by [23], with specific synthesized primers for the following resistance genes:

BLA TEM F- ATGAGTATTCAACATTTCCG
R- CTGACAGTTACCAATGCTTA

BLA SHV F- CGCCTGTGTATTATCTCCCTGTTAGCC
R- TTGCCAGTGCTCGATCAGCG

CTX-M F- CGCTTTGCGATGTGCAG
R- ACCGCGATATCGTTGGT

From Inqaba Biotec African Genomics Company. The final amplified products were electrophoresed through agarose gels (1%) containing 0.5% ethidium bromide and visualized under UV transilluminator Documentation unit [23].

Antimicrobial Assay

The antimicrobial activity of selected extracts was determined using the disc diffusion method according to [24]. Mueller Hinton agar was sterilized by autoclaving at 121°C for 15min, cooled, poured into Petri dishes and inoculated with the selected isolates by striking the swab over the

surface of the medium in three directions to confirm a complete distribution. Sterile filter paper discs (Whatman No. 3, 6 mm diameter and three layers) was saturated by stock solutions of 100µL of each extract (10 mg/mL); the disks were allowed to dry for one hour, then placed on the surface of inoculated plates. The used organic solvents and distilled water disks served as negative controls. The plates were kept in a refrigerator for one hour to allow better diffusion of the extract prior to incubation at 37°C/24h. After incubation, the inhibition zones formed around disks were measured in millimeters (including the diameter of the disk (6 mm)). Each experiment was run in triplicates and the means calculated.

In-Vivo Antimicrobial Assay

All the applicable guidelines for the care and use of animals were followed in the study. Seven days prophylactic and 1h post infection effects of *P. aeruginosa* and plants extracts was investigated using the method previously described by [25].

Statistical Analysis

Results obtained were presented as mean $\pm$ standard deviation of three determinations. Means were also analyzed using one-way analysis of variance, followed by the Duncan Multiple Range Test to determine significant differences in all the parameters using SPSS. Differences with values of $P < 0.05$ will be considered statistically significant.

Results

Plant Extraction Result

Extraction of *Jatropha curcas* and *Acacia nilotica*, chloroform extract, reveals that some of the leave extracts has a dark brownish and/or greenish colour, with a gummy or crystalline texture and different percentage yields.

Table 1: Percentage yield and physical properties of *Jatropha curcas* and *Acacia nilotica* Leave Extract

Plant Name	Initial Weight of Plant Material (g)	Total yield (g)	Yield (%)	Colour	Texture
<i>Jatropha curcas</i>	500	51.99	10.40	Dark Brownish	Crystalline
<i>Acacia nilotica</i>	500	67.03	13.41	Dark greenish	Gummy

Result for Phytochemical Analysis

Table 2-3, shows the result for qualitative and quantitative phytochemical screening of *Jatropha curcas* and *Acacia nilotica* leave extract respectively. Qualitative screening of the plants extracts reveals the presence of numerous phytometabolites, among which; tannins, saponin, alkaloids, flavonoids, phenols, terpenoids, quinones, phytosterols and

phlobatanins, are all found to be present in all the extracts. However phytochemicals like resins, chalcones, vitamin D, acidic compounds and anthoquinones are only found in some of the extracts. Quantitative phytochemical analyses of the plants extracts indicates that phenolics have a relatively high percentage concentration in all the extracts.

Table 2: Qualitative and Quantitative Phytochemical Content of Chloroform extract of *Jatropha curcas* Plant Parts

S/N	Phytochemical	Leave Extract (mg/g dry wt)	
		Qualitative	Quantitative
1	Flavonoid	+	4.05 $\pm$ 1.28
2	Alkaloid	+	9.27 $\pm$ 0.70
3	Saponins	+	2.90 $\pm$ 2.92
4	Phytosterols	+	0.63 $\pm$ 0.73
5	Phenols	+	9.55 $\pm$ 2.35
6	Terpenoids	+	1.68 $\pm$ 0.94
8	Triterpenoids	+	
9	Tannins	+	2.53 $\pm$ 1.36
10	Cardiac glycoside	+	0.18 $\pm$ 0.18
11	Anthraquinones	+	0.59 $\pm$ 1.77
12	Anthocyanins	—	
13	Phlobatanins	+	
14	Flavonols/flavones	+	
15	Coumarins	—	

16	Quinones	+	
17	Resins	+	
18	Amino acids	—	
19	Chalcones	—	
20	Vitamin A	—	
21	Vitamin D	—	
22	Acidic compound	+	

Key:

+ = Presence - = Absence

Results are presented as mean $\pm$ standard deviation

Table 3: Qualitative and Quantitative Phytochemical Content of Chloroform extract of *Acacia nilotica* Plant Parts

S/N	Phytochemical	Leave Extract (mg/g dry wt)	
		Qualitative	Quantitative
1	Flavonoid	+	1.97 $\pm$ 1.64
2	Alkaloid	+	1.53 $\pm$ 2.85
3	Saponins	+	1.32 $\pm$ 2.75
4	Phytosterols	+	0.29 $\pm$ 1.64
5	Phenols	+	9.92 $\pm$ 1.15
6	Terpenoids	+	1.32 $\pm$ 1.74
8	Triterpenoids	—	
9	Tannins	+	1.36 $\pm$ 1.82
10	Cardiac glycoside	+	0.83 $\pm$ 1.28
11	Anthraquinones	+	0.06 $\pm$ 0.94
12	Anthocyanins	—	
13	Phlobatannins	+	
14	Flavonols/flavones	+	
15	Coumarins	—	
16	Quinones	+	
17	Resins	—	
18	Amino acids	—	
19	Chalcones	—	
20	Vitamin A	—	
21	Vitamin D	—	
22	Acidic compound	—	

Key:

+ = Presence - = Absence

Results are presented as mean $\pm$ standard deviation

Result for the Isolation of *Pseudomonas aeruginosa* clinical Isolates

A total of 108 clinical samples were collected and processed from different general hospitals in Kaduna state metropolis, of which 43 (39.81) were identified as *Pseudomonas aeruginosa*. There was a considerable variation in the infection of *Pseudomonas aeruginosa* as classified by the


clinical sources. Wound samples however recorded the highest figures {18 (16.67)}, while the least was found in stool and sputum (Table 4). Table 5, however shows the socio-demographic characteristics of all the participating patients, their gender, age and a statistics of the bacterial isolation from the different groups.

Table 4: *Pseudomonas aeruginosa* clinical isolates from different General Hospitals in Kaduna metropolis

Sources (s)	No of Samples	<i>P. aeruginosa</i> Isolates (%)
Wound	38	18 (16.67)
Stool	14	3 (2.78)
Oral	7	5 (4.62)
Fluids	9	5 (4.62)
Ear Swap	11	6 (5.56)
Urine	21	6 (5.56)
Sputum	8	- (0)
Total	108	43 (39.81)

Table 5: Socio-demographic Characteristics of all the Participating Patients for the Clinical Isolation *P. aeruginosa*

Variable (s)	No of Samples (%)	No. of <i>P. aeruginosa</i> Isolates (%)
Gender		
Male	74 (68.52)	29 (26.85)
Female	34 (31.48)	14 (12.96)
Age (Range)		
≤ 20	49 (45.37)	18 (16.67)
21 – 30	22 (20.37)	13 (12.04)
31 – 40	18 (16.67)	8 (7.41)
41 – 50	8 (7.41)	1 (0.93)
50 +	11 (10.19)	3 (2.78)

**Fig 1:** *Pseudomonas aeruginosa* Isolation from Clinical Samples


Result for the Isolation of *Pseudomonas aeruginosa* Environmental Isolates

A total of 42 environmental samples were collected and processed from different parts of Kaduna state metropolis, of

which 22 (50.00) were identified as *Pseudomonas aeruginosa*. Refuse Dump samples recorded the highest figures of 7 (16.67), while the least was found in Hospitals Surgical Equipment's (Table 6).

Table 6: *Pseudomonas aeruginosa* Environmental isolates from different parts of Kaduna metropolis

Sources (s)	No of Samples	<i>P. aeruginosa</i> Isolates (%)
Refuse Dump	16	7 (16.67)
Hospital Refuse Dump	10	6 (14.28)
Poultry Houses	6	6 (14.28)
NNPC Waste Water Outlet	2	1 (2.38)
Mechanic Workshop	5	2 (4.76)
Hospitals Surgical Equipment's	3	0 (0)
Total	42	22 (50.00)

**Fig 2:** *Pseudomonas aeruginosa* Isolation from different Environmental Samples

Result for the Isolation of *Staphylococcus aureus* Clinical Isolates

A total of 89 clinical samples were collected and processed from different general hospitals in Kaduna state metropolis, of which 43 (48.31%) were identified as *Staphylococcus aureus*. There was a considerable variation in the infection of *Staphylococcus aureus* as classified by the clinical sources.

Wound samples however recorded the highest figures {13 (14.61)}, while the least was found in sputum sample (Table 7). Table 8, however shows the socio-demographic characteristics of all the participating patients, their gender, age and a statistics of the *S. aureus* isolation from the different groups.

Table 7: *Staphylococcus aureus* clinical isolates from different General Hospitals in Kaduna metropolis

Sources (s)	No of Samples	<i>Staphylococcus aureus</i> (%)
Wound	21	13 (14.61)
Stool	11	6 (6.74)
Oral	8	4 (4.49)
Fluids	9	6 (6.74)
Ear Swap	8	4 (4.49)
Urine	22	8 (8.99)
Sputum	10	2 (2.25)
Total	89	43 (48.31)

Table 8: Socio-demographic Characteristics of all the Participating Patients for the Clinical Isolation *S. aureus*

Variable (s)	No of Samples (%)	No. of <i>S. aureus</i> Isolates (%)
Gender		
Male	31 (34.83)	19 (21.35)
Female	58 (65.17)	24 (26.97)
Age (Range)		
≤ 20	18 (20.22)	10 (11.24)
21 – 30	33 (37.08)	12 (13.48)
31 – 40	30 (33.71)	17 (19.10)
41 – 50	6 (6.74)	3 (3.37)
50 +	2 (2.25)	1 (1.12)




Fig 3: *Staphylococcus aureus* Isolation from Clinical Samples

Result for the Isolation of *Staphylococcus aureus* Environmental Isolates

A total of 61 environmental samples were collected and processed from different parts of Kaduna state metropolis, of

which 44 (72.13%) were identified as *Staphylococcus aureus*. Refuse Dump samples recorded the highest figures of 24 (39.34%), while the least was found in Hospitals Surgical Equipment's (2 {3.28%}) (Table 9).

Table 9: *Staphylococcus aureus* Environmental isolates from different parts of Kaduna metropolis

Sources (s)	No of Samples	<i>Staphylococcus aureus</i> (%)
Refuse Dump	28	24 (39.34)
Hospital Refuse Dump	14	4 (6.56)
Poultry Houses	7	7 (11.48)
NNPC Waste Water Outlet	4	4 (6.56)
Mechanic Workshop	3	3 (4.92)
Hospitals Surgical Equipment's	5	2 (3.28)
Total	61	44 (72.13)




Fig 4: *Staphylococcus aureus* Isolation from different Environmental Samples

Bioassay Result for the most Resistant Strains of *Pseudomonas aeruginosa*

Antimicrobial activity assay done on all the clinical and environmental isolates done on all the 150 collected samples,

reveals that eleven of the isolates are extreme multi-drug resistant, based on the bioassay analyses carried out using standard antibiotics (Table 10).

Table 10: Bioassay for the most resistant strains of *P aeruginosa*

Resistant Isolates	Standard Antibiotics Zone of Inhibition in cm												
	NF	CXM	CRO	ACX	ZEM	LBC	AUG	CTX	IMP	OFX	GN	NA	VA
P ₃	-	-	-	-	-	1.9	-	-	-	-	-	0.9	-
P ₁₅	-	-	-	-	-	-	1.3	-	-	1.4	-	-	-
P ₁₆	-	-	-	-	-	-	-	-	1.2	-	-	-	-
P ₂₀	-	-	1.4	-	-	-	-	-	-	-	-	-	-
P ₂₆	-	-	-	-	-	1.0	-	-	-	1.5	-	-	-
P ₄₂	-	-	-	-	-	-	-	-	0.9	-	1.6	-	-
P ₄₇	-	1.2	-	-	-	-	1.1	-	-	-	-	-	-
P ₈₈	-	-	-	-	-	0.9	-	-	-	-	1.0	-	-
P ₁₀₀	-	-	-	-	-	-	-	-	-	1.2	1.1	-	-
P ₁₂₅	-	-	-	-	-	-	-	-	-	-	-	-	1.4
P ₁₄₆	-	-	-	-	-	1.7	-	-	-	-	1.4	-	-

Key: P: *P aeruginosa*, NF - Nitrofurantoin (300µg), CXM – Cefuroxime (30µg), CRO – Ceftriaxone Sulbactam (45µg), ACX - Ampiclox (10µg), ZEM - Cefexime (5µg), LBC – Levofloxacin (5µg), AUG - Amoxicilin Clavulanate (30µg), CTX – Cefotaxime (25µg), IMP - Imipenem/Cilastatin (10/10µg), OFX - Ofloxacin (5µg), GN – Gentamycin (10µg), NA - Nalidixic Acid (30µg) and VA – Vancomycin (30µg)

Bioassay Result for the most Resistant Strains of *Staphylococcus aureus*

Antimicrobial activity assay done on all the clinical and environmental *S. aureus* isolates done on all the 150 collected

samples, reveals that four of the isolates are multi-drug resistant, based on the bioassay analyses carried out using standard antibiotics (Table 11).

Table 11: Bioassay for the most resistant strains of *S. aureus*

Resistant Isolates	Standard Antibiotics Zone of Inhibition in cm											
	PEF	CN	APX	Z	AM	R	CPX	S	SXT	E	M	
S ₃₇	-	1.6	-	-	-	-	-	-	-	-	-	-
S ₆₄	-	-	-	-	-	-	1.0	1.2	-	-	-	-
S ₇₂	1.4	-	-	-	-	-	-	-	-	0.8	-	-
S ₁₀₅	-	-	-	1.4	-	-	-	-	-	-	-	-

Key: PEF - Pefloxacin (10µg), CN - Gentamycin (10µg), APX - Ampiclox (30µg), Z - Zinnacef (20µg), AM - Amoxicillin (30µg), R - Rocephin (25µg), CPX - Ciprofloxacin (10µg), S - Streptomycin (30µg), SXT - Septrin (30µg), E – Erythromycin (10µg) and M – Methicillin (30µg)

Results for the Resistance Genes Identification

The agarose Gel Electrophoresis of the PCR amplified products of Extended-Spectrum Beta-lactamase Genes done,

revealed the presence of the selected resistance genes in virtually all the selected extremely multi-drug resistant bacterial strains (Figure 5).




Fig 5: Agarose gel electrophoresis of CTX-M, bla SHV and bla TEM (multiplex PCR), M: BIONEER DNA Marker 100bp

Results for the Susceptibility Antibigram Profile of the Two Bacteria to Different Antibiotic Groups

Susceptibility pattern analyses reveals that the two bacterial

resistant strains have different individual susceptibility and resistance to various antibiotic groups (Table 12 & 13).

Table 12: *Pseudomonas aeruginosa* Susceptibility Pattern to Different Antibiotics Groups (n=150)

S/N	Antibiotics Group	No. of Susceptible	No. of Resistant
1	Cephalosporins	347	253
2	Penicillins	171	129
3	Quinolones	267	183
4	Carbapenems	101	49
5	Aminoglycosides	88	62
6	Glycopeptides	77	73
7	Nitrofurantoin	81	69

Table 13: *Staphylococcus aureus* Susceptibility Pattern to Different Antibiotics Groups (n=150)

S/N	Antibiotics Group	No. of Susceptible	No. of Resistant
1	Cephalosporins	196	104
2	Penicillins	277	173
3	Quinolones	191	109
4	Macrolides	105	45
5	Aminoglycosides	200	100
6	Sulfonamides and Trimethoprim	96	54

In-vitro Bioassay of Plants Extracts against Multidrug Resistant Isolates of *Pseudomonas aeruginosa*

Mean Results for the In-vitro bioassay of *J. curcas* and *A. nilotica* leave extracts against multidrug resistant isolates of

Pseudomonas aeruginosa, which reveals that *J. curcas* extract has the highest antimicrobial activity against the various multi-drug resistant strains (Table 14).

Table 14: Plants extracts bioassay against multidrug resistant isolates of *Pseudomonas aeruginosa*

S/N	Resistant Isolates	Isolate Source	Crude Extracts Zone of Inhibition (cm) (1000µg/ml)		Control (cm)
			<i>Jatropha curcas</i>	<i>Acacia nilotica</i>	Gentamycin
1	P ₃	Clinical	1.3 ± 2.04	1.6 ± 1.05	-
2	P ₁₅	Clinical	1.2 ± 1.18	1.5 ± 1.72	-
3	P ₁₆	Clinical	1.5 ± 0.21	1.1 ± 1.49	-
4	P ₂₀	Env.	0.6 ± 1.05	1.0 ± 0.56	-
5	P ₂₆	Clinical	2.0 ± 0.44	0.6 ± 1.43	0.5
6	P ₄₂	Clinical	1.6 ± 1.10	1.7 ± 1.01	1.4
7	P ₄₇	Clinical	1.8 ± 0.04	2.7 ± 1.47	-

8	P ₈₈	Clinical	1.4 ± 1.62	1.9 ± 0.05	0.8
9	P ₁₀₀	Clinical	2.3 ± 1.03	1.7 ± 0.53	1.1
10	P ₁₂₅	Env.	1.4 ± 0.74	1.3 ± 1.46	-
11	P ₁₄₆	Clinical	2.0 ± 1.02	0.9 ± 0.85	1.3

Key:

P: *Pseudomonas aeruginosa*

±: Standard Deviation

-: No zone of inhibition

Env: Environmental

In-vitro Bioassay of Plants Extracts against Multidrug Resistant Isolates of *Staphylococcus aureus*

Mean Results for the in-vitro bioassay of *J. curcas* and *A. nilotica* leave extracts against multidrug resistant isolates of

Staphylococcus aureus in cm. The analysis reveals that *J. curcas* has the highest antimicrobial activity against the different strains of multi-drug resistant *S. aureus* (Table 15).

Table 15: Plants extracts bioassay against multidrug resistant isolates of *S. aureus*

S/N	Resistant Isolates	Isolate Source	Crude Extracts Zone of Inhibition (cm) (1000µg/ml)		Control (cm)
			<i>Jatropha curcas</i>	<i>Acacia nilotica</i>	Gentamycin
1	S ₃₇	Clinical	2.1 ± 1.11	1.9 ± 0.18	1.5
2	S ₆₄	Clinical	1.2 ± 1.80	0.7 ± 0.56	-
3	S ₇₂	Env.	1.9 ± 0.83	1.9 ± 1.03	-
4	S ₁₀₅	Clinical	2.1 ± 2.07	0.9 ± 0.55	-

Key:

S: *Staphylococcus aureus*

±: Standard Deviation

-: No zone of inhibition

Env.: Environmental

In-vitro Bioassay of Decocted Extract of *J. curcas* and *A. nilotica* against Extreme Multidrug Resistant Isolates of *Pseudomonas aeruginosa* and *Staphylococcus aureus*

Mean Results for the in-vitro bioassay of a decocted extract, against the isolated multidrug resistant isolates of

Pseudomonas aeruginosa, shows that combining the three most bio-active extracts proves to be more efficient in inhibiting the growth of the extreme multi-drug resistant strains of both the virulent bacteria (Table 16).

Table 16: Extracts decoction of *J. curcas* and *A. nilotica* bioassay against extreme multidrug resistant isolates of *Pseudomonas aeruginosa* and *Staphylococcus aureus*

S/N	Resistant Isolates	Isolate Source	Decocted Extract Zone of Inhibition (cm) (1000µg/ml)	Control Gentamycin (cm)
1	P ₃	Clinical	2.6 ± 1.53	-
2	P ₁₅	Clinical	2.8 ± 0.64	-
3	P ₁₆	Clinical	2.4 ± 1.32	-
4	P ₂₀	Env.	1.3 ± 1.03	-
5	P ₂₆	Clinical	1.4 ± 0.24	1.0
6	P ₄₂	Clinical	2.5 ± 2.72	1.1
7	P ₄₇	Clinical	1.1 ± 1.29	-
8	P ₈₈	Clinical	2.0 ± 0.34	0.5
9	P ₁₀₀	Clinical	2.6 ± 1.35	1.2
10	P ₁₂₅	Env.	1.7 ± 2.31	-
11	P ₁₄₆	Clinical	2.4 ± 1.07	1.0
12	S ₃₇	Clinical	2.9 ± 2.05	1.2
13	S ₆₄	Clinical	2.3 ± 1.01	-
14	S ₇₂	Env.	2.8 ± 1.15	-
15	S ₁₀₅	Clinical	3.1 ± 1.78	-

Key:

P: *Pseudomonas aeruginosa*

S: *Staphylococcus aureus*

±: Standard Deviation

-: No zone of inhibition

Env.: Environmental

Results for the In-vivo Bioassay of the Plants Extracts against Resistant Isolates of *Pseudomonas aeruginosa*

The in-vivo bioassay of the two plants leave extracts against multidrug resistant isolates of *Pseudomonas aeruginosa* in

Rats Model (3 Rats per Group unit) reveals that both *J. curcas* and *A. nilotica* extracts are quite very active against the activity of the resistant strains of *P. aeruginosa* (Table 17).

Table 17: In-vivo bioassay of the plants extracts against *P. aeruginosa* in rats model

S/N	Resistant Isolates	Isolate Source	Infected Groups + Crude Extracts (500µg/ml)		Control	
			<i>Jatropha curcas</i> (33)	<i>Acacia nilotica</i> (33)	+	-
1	P ₃	Clinical	3	2	3	0
2	P ₁₅	Clinical	2	3		0
3	P ₁₆	Clinical	1	3		0
4	P ₂₀	Env.	1	2		0
5	P ₂₆	Clinical	3	2		0
6	P ₄₂	Clinical	3	2		1
7	P ₄₇	Clinical	3	3		0
8	P ₈₈	Clinical	2	3		0
9	P ₁₀₀	Clinical	3	3		1
10	P ₁₂₅	Env.	0	2		0
11	P ₁₄₆	Clinical	2	1		0
Total Remaining			23	26	3	2

Key:

P: *Pseudomonas aeruginosa*

Env: Environmental

+: Positive Control (Un-infected)

-: Negative Control (Infected)

Results for the In-vivo Bioassay of the Plants Extracts against Resistant Isolates of *S. aureus*

The in-vivo bioassay of plants leave extracts against multidrug resistant isolates of *S. aureus* in Rats Model (3 Rats

per Group unit) reveals that *J. curcas* extracts is the most bioactive bio-metabolite against the activity of the resistant strains of *S. aureus* (Table 18).

Table 18: In-vivo bioassay of the plants extracts against *S. aureus* in rats model

S/N	Resistant Isolates	Isolate Source	Infected Groups + Crude Extracts (500µg/ml)		Control	
			<i>Jatropha curcas</i> (12)	<i>Acacia nilotica</i> (12)	+	-
1	S ₃₇	Clinical	3	2	3	0
2	S ₆₄	Clinical	3	1		0
3	S ₇₂	Env.	2	3		1
4	S ₁₀₅	Clinical	2	3		0
Total (Remaining)			10	9	3	1

Key:

S: *Staphylococcus aureus*

Env.: Environmental

+: Positive Control (Un-infected)

-: Negative Control (Infected)

Discussion

Medicinal plants has for quite a while use as potential source of pharmaceutical agents and traditionally used in the treatment and management different kinds of diseases and infections, such as diarrhea, fever and cold [1]. These plants parts, decoction, phyto-metabolites (which are metabolic chemical compounds that occur naturally in plants [26], responsible for a lot of plants peculiar features and capability, such as; medical defense, color and organoleptic properties) and other matter are nowadays, widely used in conventional and alternative medical practices in developed and developing countries as a complementary medicine [2]. However emergence of new strains of pathogenic agents, couple with the detrimental effects of multi-drug resistance, especially among gram negative and positive bacteria, which now adays always increasing by the day, making even some of the most effective drugs and antibiotics available useless. Resistant bacterial strains like *P. aeruginosa* and *S. aureus* can use both intrinsic, acquired and adaptive microbial antibiotic resistance mechanisms to rapidly develop resistance to various kinds of bacterial antibiotics, such as the cephalosporin's, aminoglycosides, quinolones and β-lactams. Plants phytometabolites can however be exploits, in the treatment and management of such ailments cause by multi-drugs resistant pathogens, which other available

pharmaceutics have failed to manage [27]. In this research antibacterial activity of five selected ethno-medicinal plants have been research, against some of the most multi-drug resistant strains of *Pseudomonas aeruginosa* and *Staphylococcus aureus*, isolated from various general hospitals, clinical environment and different kinds of environmental focal points in Kaduna state metropolis, Nigeria. The plant extraction was done using chloroform as the extraction solvent, in which *Acacia nilotica* plant material extraction yielded the highest percentage yield of 13.41% (Table 1). Qualitative phytochemical screening of the plants extracts done, shows that all the five plants contain; tannins, saponin, alkaloids, flavonoids, phenols, terpenoids, quinones, phytosterols and phlobatanins in them. While quantitative phytochemical analyses reveals that all the extracts contains phenolic chemicals as the highest containing chemicals in them, however *F. sycomorus* and *J. curcas* extracts appears to be the two most chemically enriched extracts among the five selected plants. Some of these findings were similarly reported by researchers like; [16, 28, 29, 30, 31, 32, 33, 34].

One hundred and eight (108) clinical samples were collected from different general hospitals for the isolation of *P. aeruginosa*, out of which 39.81% were found to *P. aeruginosa* strains, of which most the strains are of Wound swap samples (Figure 1). While isolation of same strains

from environmental origins was also done using forty two (42) collected samples of *P. aeruginosa* strains were successfully isolated from 50% of the sample. Refuse dump samples however gave the highest percentage isolates from the various samples collected (Figure 2). On the other hand Eighty nine (89) clinical samples were collected from different general hospitals also for the isolation of *S. aureus*, out of which 48.31% were found to be positive for *S. aureus* presence (Figure 3), of which most the strains are also of wound swap origin, just like that of *P. aeruginosa* isolation. While isolation of same strains from environmental origins was also done using 61 collected samples of *S. aureus* strains were successfully isolated from 72.13% of the samples. Refuse dump samples however also gave the highest percentage isolates from the various environmental samples collected (Figure 4).

Standard antimicrobial bioassay done on all the clinical and environmental *P. aeruginosa* isolates, reveals that eleven of the isolates are multi-drug resistant, based on the bioassay analyses carried out using gram negative standard antibiotic assay disc. Eleven clinical and environmental *P. aeruginosa* strains were found to be extremely multi-drug resistant to more than ten (10) standard antibiotics, including drugs of choice for the bacteria (Imipenem/Cilastatin). While bioassay sensitivity analyses done on *S. aureus* isolates reveals that four (4) of the 150 isolated strains are also extremely multi-drug resistant to numerous antibiotics, including methicillin. Of which clinical samples; S₃₇ and S₁₀₅ are the most resistant isolates, which are only susceptible to gentamycin and Zinnacef respectively.

Resistance genes identification analyses done reveals that most of the isolated resistance strains have either two or three of the BLA-TEM, BLA-SHV and CTX-M resistance genes (Figure 5). These genes have been reported to be responsible for multiple antibiotic drug resistance mechanisms including protein activators for multiple efflux pump systems and quorum sensing activation and enhancement [35].

In-vitro analyses done on the multi-drug resistant isolates against the two selected plants leave extract shows that both the plants have a relatively high antimicrobial effects against most of the resistant strains, however *J. curcas* extracts has the highest antimicrobial activity against the various multi-drug resistant strains. The antibiotic gentamycin was used as a reference control drug, as it was initially used on both the gram positive and negative isolates during the anti-biogram analyses. This findings were similar with the research done by [36], on biodiesel industrial waste based on *Jatropha curcas* as a fungicide to control *Fusarium oxysporum* and *Alternaria solani*. The analyses also shows that *P. aeruginosa* strain 'P₂₀', which is an environmental isolate from a mechanic workshop, is the most dominantly multi resistant strain of bacteria among all the selected resistant isolates. The resistant strains in this research are quite aggressively resistant to almost all kinds of antibiotics including the drug of choice for the multi-drug resistant isolates. For example in contrast to a research done by [37], which reported that most extremely resistant strains of *P. aeruginosa* and *S. aureus* were susceptible to Vancomycin and Imipenem, which is not in line with these isolated strains of multi-drug resistant bacteria. This aggressive resistibility might be as a result of an acquired mutation from multiple contacts with some of the various chemical compounds from the mechanic workshop, couple with the *P. aeruginosa* natural multi-drug resistance genes and intrinsic factors. Decoction of the two plants

extracts (*J. curcas* and *A. nilotica*), proof effective, revealing a relatively high synergistic effect of the extract to each other. In-vivo antimicrobial bioassay done, using albino rats shows the pathogenic infectivity of all the resistance strains, as the isolates were able to kill all the infected groups after one days post infection study. *Acacia nilotica* extracts however recorded the highest in-vivo anti-bacterial effects on the various resistance strains, than the *J. curcas* extracts, which activity has already been established from the in-vitro bioassay done. This research is also compare with the research done by [28] on the phytochemical properties, toxicological screening and antibacterial qualities of various parts extracts of *Ficus sycomorus* and the research done by [15] on the Antimicrobial Activities of *Cymbopogon citratus* and *Ximenia Americana* Leaf Extracts Against Some Selected Bacterial and Yeast Clinical Isolates.

Conclusion

J. curcas and *A. nilotica* plant extracts could provide an alternative natural pharmaceutical agent or semi decoction part of a drug formulation for the management and treatment of various kinds of ailments cause by multi-drug resistance *P. aeruginosa* and *S. aureus* bacterial pathogens.

Reference

1. Hassan AU, NE Egbe, J Appah, S Garba a, Plants Phenoliscs as Possible *Pseudomonas aeruginosa* Resistance Inhibitors. Asian Plant Research Journal. 2023; 11(1):1-9. DOI: 10.9734/APRJ/2023/v11i1200.
2. Abu MS, OE Yakubu, RN Boyi, MH Mayel, CO Shaibu. Phytochemical constituents, acute toxicity and free radical scavenging activity of methanol extract of *F. glumosa* leaves. Anchor Univ J Sci Technol. 2020; 1(1):16-22. <https://www.ajol.info/index.php/aujst/article/view/222125>
3. Umaru IJ, OE Yakubu, MU Ahmed, I Ezekiel, KI Umaru, MS Abu, M Abah. GC-MS, radicals scavenging capacity and antidiabetic effect of senna alata seed extract in type ii-induced diabetes mellitus in rats. Research Journal of Medicinal Plants. 2022; 16(2):49-58. 2022. <https://doi.org/10.17311/rjmp.2022.49.58>
4. Rivas MÁ, Casquete R, de Guía Córdoba M, Benito MJ, Hernández A, Ruiz-Moyano S, et al. Functional properties of extracts and residual dietary fibre from pomegranate (*Punica granatum* L.) peel obtained with different supercritical fluid conditions. LWT-Food Sc. Tech. 2021; 145:111305
5. Olanipekun MK, K Akirun, JA Amoo. Phytochemical Screening and Proximate Analyses of Some Medicinal Plants Used in Irun Akoko, Akoko North West Local Government Area Ondo State, Nigeria (July 12, 2021). European Journal of Botany, Plant Sciences and Phytology. 2021; 6(1):1-13.
6. Breidenstein EBM, De la Fuente-Núñez C, Hancock REW. *Pseudomonas aeruginosa*: all roads lead to resistance. Trends in Microbiology A/I press Cambridge. 2011; 19:419-426.
7. Ejaz H, Younas S, Abosalif KOA, Junaid K, Alzahrani B, Alsrhani A, et al. Molecular analysis of blaSHV, blaTEM, and blaCTX-M in extended-spectrum β -lactamase producing Enterobacteriaceae recovered from fecal specimens of animals. PLoS ONE. 2021; 16(1):e0245126. doi:10.1371/journal.pone.0245126

8. Ehlers Marthie M, Chrisna Veldsman, Eddy P. Makgotlho, Michael G. Dove, Anwar A. Hoosen and Marleen M. Kock Detection of blaSHV, blaTEM and blaCTX-M antibiotic resistance genes in randomly selected bacterial pathogens from the Steve Biko Academic Hospital. FEMS Immunol Med Microbiol. 2009; 56:191-196.
9. Shikov AN, Shikova VA, Whaley AO, Burakova MA, Flisyuk EV, Whaley AK, Terninko II, *et al.* The Ability of Acid-Based Natural Deep Eutectic Solvents to Co-Extract Elements from the Roots of Glycyrrhiza glabra L. and Associated Health Risks. Molecules. 2022; 27:7690.
10. Pecina Quintero V, Anaya-López JL, Zamarripa-Colmenero A, Núñez-Colín CA. Genetic structure of *Jatropha curcas* L. in Mexico and probable centre of origin. Biomass Energy. 2014; 60:147-155 <http://dx.doi.org/10.1016/j.biombioe.2013.11.005>.
11. Gamal Fakhry, Azazz NAE, Abdel-Monem A, Mohamedin A. Use of waste water and treated water for *jatropha curcas* cultivation and the possibility of oil seed use as a biofuel. Minia J of Agric. Res. & Develop. 2016; 36(2):245-269.
12. Kalaivani T, Mathew. Free radical scavenging activity from leaves of *Acacia nilotica* (L.) Wil. Ex-Delile, an Indian medicinal tree. Food Chem. Toxicology. 2010; 48:298-305.
13. Jigam AA, Akanya HO, Dauda BEN, Okogun JO. Polygalloyltannin isolated from the roots of *Acacia nilotica* Del. (Leguminosae) is effective against *Plasmodium berghei* in mice. J Med. Plants Res. 2010; 4(12):1169-1175.
14. Aliyu B S. (2006): Common Ethnomedicinal Plants of the Semiarid Regions of West Africa; Triumph Publishing Company Limited; Nigeria Pp 8, 26.
15. Hassan A. U., A. H. Madu, U. O. Ozojiofor, A. H. Galadanci, I. B. Mato and R. Jafaru (2021), Antimicrobial Activities of *Cymbopogon citratus* and *Ximenia Americana* Leaf Extracts Against Some Selected Bacterial and Yeast Clinical Isolates. *Asian Journal of Biochemistry, Genetics and Molecular Biology*; 9(1): 1-10, DOI: 10.9734/AJBGMB/2021/v9i130204
16. Hassan AU, MS Makeri, UO Ozojiofor, AJ Alhassan. Effects of *Ficus sycomorus* Phenolic Extracts on the Activity of Hyaluronidase and Phospholipase A₂ Enzymes of *Echis ocellatus* Venom. *Asian Journal of Biotechnology and Genetic Engineering*. 2020; 3(4):24-33.
17. Lakache Z, Tigrine-Kordjani N, Tigrine C, Aliboudhar H, Kameli A. Phytochemical screening and antioxidant properties of methanolic extract and different fractions of *Crataegus azarolus* leaves and flowers from Algeria. *International Food Research Journal*. 2016; 23(4):1576-1583
18. Gellatly SL, Hancock R. *Pseudomonas aeruginosa*: New insights into pathogenesis and host defenses. *Pathogens and Disease*. 2013; 67(3):159-173.
19. Brown A, Horobin A, Blount DG, Hill PJ, English J, Rich A, Pritchard DI. Blow fly *Lucilia sericata* nuclease digests DNA associated with wound slough/eschar and with *Pseudomonas aeruginosa* biofilm. *Medical and veterinary entomology*. 2012; 26(4):432-439
20. Sarwat T, Rashid M, Rastogi V, Chander Y. Original research article a comparative study of antibiogram of *Pseudomonas aeruginosa* in hospital and community acquired infections. *International Journal of Curriculum Microbiology Application Science*. 2015; 1:286-291.
21. Kossah R, Nsabimana C, Zhang H, Chen W. Evaluation of antimicrobial and antioxidant activities of Syrian Sumac fruit extract. *J Nat. Prod*. 2013; 6:96-102.
22. Cheesbrough M. *District Laboratory Practice in Tropical Countries*. Published by press syndicate of the University of Cambridge, the Edinburg building, Cambridge United Kingdom, NewYork, 2000, 194-201.
23. Shaebth LJ. Molecular identification and sequencing of *Pseudomonas aeruginosa* virulence genes among different isolates in Al-Diwaneyah hospital. *Iraqi Journal of Veterinary Sciences*. 2018; 32(2):183-188.
24. Black JG, Black LJ. *Microbiology: Principles and Explorations*; John Wiley & Sons: Hoboken, NJ, USA, 2018.
25. Oke AJ. Investigation of Immunomodulatory properties of Protein A from a Local strain of *Staphylococcus aureus*, Ph. D thesis Dissertation, Ladoke Akintola University of Technology, Ogbomoso, 2006.
26. Gong T, Liu S, Wang H, Zhang M. Supercritical CO₂ fluid extraction, physicochemical properties, antioxidant activities and hypoglycemic activity of polysaccharides derived from fallen Ginkgo leaves. *Food Biosci*. 2021; 42:101153.
27. Michael Sunday Abu, Rukaiyat Lawal Mashi, Job Itanyi Onuche and Sa'ad Ismail Shuaibu. Proximate and phytochemical screening of some selected herbs and spices commonly used in Nigeria. *GSC Biological and Pharmaceutical Sciences*. 2023; 22(01):015-024
28. Braide W, Dokubo KO, Adeleye SA, Uzoh CV, Akobundu CI. Phytochemical properties, toxicological screening and antibacterial qualities of various parts extracts of *Ficus sycomorus*. *J Med Plant Herb Ther Res*. 2018; 6:1-8.
29. Hassan AU, Girbobo BU, Umar SI. b, Fungal load evaluation of the leave and fruit extracts of *Vitex Doniana* (black plum) using three basic phyto-medicinal preparatory protocols. *International Journal of Medical and All Body Health Research*. 2023; 04(01):05-10.
30. Bunu MI, Ndinteh DT, Macdonald JR, Langat MK, Isyaka SM, Sadgrove NJ, *et al.* Ecdysteroids from the Stem Bark of *Vitex doniana* Sweet (Lamiaceae; ex. Verbenaceae): A Geographically Variable African Medicinal Species. *Antibiotics*. 2021; 10:937. <https://doi.org/10.3390/antibiotics10080937>
31. Makholwa K, Moseki B, Malambane G. Effect of Water Deficit Stress on Photosynthetic Characteristics of *Jatropha curcas*. *American Journal of Plant Sciences*. 2023; 14:104-112. <https://doi.org/10.4236/ajps.2023.141008>
32. Perumal Vivekanandhan, Swathy Kannana, Sarayut Pittaratea, Ragavendran Chinnasamyc, Patcharin Krutmuang Essential oils from *Acacia nilotica* (Fabales: Fabaceae) seeds: May have insecticidal effects?. *Heliyon*. 2023; 9:e14808. doi.org/10.1016/j.heliyon.2023.e14808
33. Alagbe JO. *Prosopis africana* (African mesquite) oil as an alternative to antibiotic feed additives on broiler chickens diets: haematology and serum biochemical indices. *Central Asian Journal of Theoretical and Applied Sciences*. 2022; 3(2):19-29.

34. Anil Kumar A, Patil NS, Kumar R, Mandal D. Irrigation Scheduling and Fertilization Improves Production Potential of *Jatropha* (*Jatropha curcas* L.). Int. J Curr. Microbiol. App. Sci. 2017; 6(5):1703-1716.
35. Nilsson LE. Multiplex PCR amplification assay for the detection of blaSHV, blaTEM and blaCTX-M genes in Enterobacteriaceae. APMIS. 2007; 115:1400-1408. doi: 10.1111/j.1600-0463.2007.00722.x.
36. Henik Sukorini1, Dyah Erni Widyastuti, Dini Kurniawati2, Sawita Suwannarat3, Maizirwan Mel4,5, and Roy Hendroko Setyobudi, Biodiesel Industrial Waste based on *Jatropha curcas* as a Fungicide to Control *Fusarium oxysporum* and *Alternaria solani*. E3S Web of Conferences. 2023, **374**, <https://doi.org/10.1051/e3sconf/202337400016>
37. Rubina Sabir, Syed Faraz, Danish Alvi, Asher Fawwad, Abdul Basit. Antibigram of *Pseudomonas aeruginosa* and Methicillin-resistant *Staphylococcus aureus* in patients with diabetes. Pak J Med Sci. 2014; 30(4): 814-18. doi: 10.12669/pjms.304.4755.