



International Journal of Medical and All Body Health Research

Assessment of Social Anxiety and Associated Factors in Patients with Cleft Lip and Palate in Ilam City in 2026

Masomeh Soltanmohamadi

Department of Psychology, Ilam Branch, Islamic Azad University, Ilam, Iran

* Corresponding Author: Masomeh Soltanmohamadi

Article Info

E-ISSN: 2582-8940

ISSN (online): 2582-8940

Volume: 07

Issue: 02

Received: 24-02-2026

Accepted: 22-03-2026

Published: 25-04-2026

Page No: 67-70

Abstract

Introduction: Cleft lip and palate are among the congenital anomalies of the head and neck that often require surgical intervention. Therefore, the present study was conducted to determine the level of social anxiety among patients with cleft lip and palate.

Methods: In this cross-sectional study, 80 patients with cleft lip and palate were evaluated. The data collection tool was the La Greca Social Anxiety Questionnaire, which consists of 16 items and measures three components. After identifying individuals with cleft lip and palate and explaining the objectives of the study, the researchers recruited participants. Finally, the collected data were entered into SPSS software version 18 and analyzed using descriptive and inferential statistical tests, and the results were reported.

Results: Result showed, 57.5% of the participants were male, 71.3% were aged 12–14 years, 73.8% lived in rural areas, 52.5% had no history of reconstructive surgery, and 97.5% were classified as having poor family financial status. The mean (standard deviation) of the total social anxiety score was 60.47 ± 6.88 . The mean score for fear of negative evaluation was 26.47 ± 5.46 , social avoidance and distress in new situations was 18.36 ± 2.08 , and general social avoidance and distress was 15.63 ± 1.75 . A significant relationship was also observed between social anxiety scores and variables such as gender, place of residence, and history of reconstructive surgery ($p < 0.05$). Female patients, those living in urban areas, and those without a history of reconstructive surgery reported higher levels of social anxiety.

Conclusion: The results showed that the mean level of social anxiety in this group was relatively high, with the highest score related to fear of negative evaluation. Social anxiety was significantly higher among females, urban residents, and individuals who had not undergone reconstructive surgery. Therefore, demographic factors and treatment status may play an important role in the severity of social anxiety in these patients.

DOI: <https://doi.org/10.54660/IJMBHR.2026.7.2.67-70>

Keywords: Social anxiety, Cleft lip and palate, Anxiety

Introduction

Anxiety, as an inseparable part of human life, involves an unpleasant and pervasive feeling of worry with an unknown origin and is often accompanied by physiological symptoms. It becomes a disorder when it is experienced subjectively, continuously, and repeatedly and interferes with individuals' daily functioning^[1,2]. Anxiety is considered the most common mental health disorder and a psychological condition that most people experience at various times in their lives. It is also the most common psychological response to changes and new experiences.

Furthermore, it is a diffuse and highly unpleasant feeling, often accompanied by one or more physical indicators such as a sensation of emptiness in the stomach, shortness of breath, palpitations, sweating, headache, urge to urinate, restlessness, and a tendency to move [3, 5].

Various definitions of anxiety have been proposed. One definition describes it as excessive fear in real or imagined situations. Anxiety disorders are characterized by chronic, excessive, and uncontrollable worry about various issues, accompanied by anxiety, hypervigilance, and related physical symptoms [5, 7]. Among the most common anxiety disorders that significantly affect an individual's daily life and functioning is Social Anxiety Disorder (SAD). SAD, also known as social phobia, refers to intense fear or anxiety in social situations in which an individual may be exposed to the attention or scrutiny of others [8, 9]. SAD is a psychological disorder characterized by intense and persistent fear of social or performance situations in which the individual fears negative evaluation by others. This disorder can lead to avoidance of social interactions, reduced occupational and academic performance, and decreased quality of life [10, 11]. The main feature of SAD is a marked fear or anxiety about social situations in which the individual may be closely observed by others. In such situations, the individual fears acting in a certain way or showing anxiety symptoms that may be negatively evaluated by others [9, 12, 13].

The body is not merely a biological and psychological entity; rather, it is also a phenomenon that is socially constructed and shaped. With the emergence of major social, cultural, and political changes in the late twentieth century, new technologies such as gene therapy, medical interventions in the body, and the growth of consumer culture in modern industrial societies have increased attention to the body and body-related behaviors in contemporary consumer societies. One of these behaviors observed in many societies is the increasing tendency toward cosmetic surgery. Among these are surgeries related to cleft lip and palate [14, 16].

Cleft lip and palate are among the congenital anomalies of the head and neck. The nature of cleft lip and palate is multifactorial, involving both environmental and genetic factors. Considering the relatively high prevalence of cleft lip and palate among children, timely and appropriate treatment of this anomaly is essential to achieve optimal outcomes [17, 18]. Therefore, the present study was conducted to determine

the level of social anxiety among patients with cleft lip and palate.

Methods

In this cross-sectional study, 80 patients with cleft lip and palate were evaluated. Before the study began, the objectives of the research were fully explained to the participants and their parents. After providing the necessary explanations and obtaining informed consent, the data collection process was initiated.

The inclusion criteria were having cleft lip and palate, being within the age range of 12 to 18 years, and willingness to participate in the study. Data collection tools included a demographic information form (age, gender, family economic status, place of residence, etc.) and a social anxiety questionnaire. The questionnaire used was the La Greca Social Anxiety Questionnaire, which consists of 16 items and measures three components: fear of negative evaluation (items 1–7), social avoidance and distress in new situations (items 8–12), and general social avoidance and distress (items 13–16). Scoring of this instrument is based on a five-point Likert scale, ranging from “very much like me” with a score of 5 to “not at all like me” with a score of 1. In this study, the total score ranged from a minimum of 16 to a maximum of 80 [19].

After identifying individuals with cleft lip and palate and providing the necessary explanations about the study objectives, the researchers recruited participants. The questionnaires were then provided to the participants and completed through interviews. Ethical considerations included obtaining informed consent, explaining the purpose of the study, and maintaining the confidentiality of participants' information.

Finally, the collected data were entered into SPSS software version 18 and analyzed using descriptive and inferential statistical tests, and the obtained results were reported.

Result

According to the findings, 57.5% of the participants were male, 71.3% were in the age range of 12–14 years, 73.8% lived in rural areas, 52.5% had no history of reconstructive surgery, and 97.5% were classified as having poor family financial status (Table 1).

Table 1: Demographic and Clinical Characteristics of Participants (n = 80)

Variable	Category	Frequency (n)	Percentage (%)
Gender	Male	46	57.5
	Female	34	42.5
Age Group	12–14 years	57	71.3
	15–18 years	23	28.7
Place of Residence	Urban	21	26.3
	Rural	59	73.8
History of Reconstructive Surgery	Yes	38	47.5
	No	42	52.5
Family Economic Status	Poor	78	97.5
	Moderate	2	2.5

According to the findings, the mean (standard deviation) of the patients' total social anxiety score was 60.47 ± 6.88 . The mean score for fear of negative evaluation was 26.47 ± 5.46 ,

for social avoidance and distress in new situations was 18.36 ± 2.08 , and for general social avoidance and distress was 15.63 ± 1.75 (Table 2).

Table 2: Central Tendency and Dispersion Indices of Social Anxiety Scores and Its Dimensions

Variable	Minimum	Maximum	Mean $\pm$ SD
Total Social Anxiety Score	45	69	60.47 $\pm$ 6.88
Fear of Negative Evaluation	16	34	26.47 $\pm$ 5.46
Social Avoidance and Distress in New Situations	10	21	18.36 $\pm$ 2.08
General Social Avoidance and Distress	11	18	15.63 $\pm$ 1.75

According to the findings, a significant relationship was observed between social anxiety scores and variables such as gender, place of residence, and history of reconstructive

surgery. Female patients, those living in urban areas, and patients without a history of reconstructive surgery reported higher levels of social anxiety (Table 3).

Table 3: Comparison of the mean social anxiety scores based on demographic and clinical variables.

Variable	Category	Mean $\pm$ SD	Test	P-value
Gender	Male	63.49 $\pm$ 5.24	t-test	0.000
	Female	53.00 $\pm$ 4.27		
Age Group	12–14 years	60.47 $\pm$ 6.92	t-test	0.999
	15–18 years	60.46 $\pm$ 6.91		
Place of Residence	Urban	65.23 $\pm$ 4.41	t-test	0.000
	Rural	58.77 $\pm$ 6.83		
History of Reconstructive Surgery	Yes	56.60 $\pm$ 6.71	t-test	0.000
	No	63.97 $\pm$ 4.92		

Discussion

Social anxiety disorder is a condition in which an individual experiences excessive and irrational fear of social situations. This disorder, as a disabling condition, has adverse effects on the quality of social relationships, motivation, academic progress, and well-being [20]. Considering the complications resulting from diseases, especially among adolescents, the present study was conducted to determine the status of social anxiety in patients with cleft lip and palate.

According to the findings, a significant relationship was observed between the level of social anxiety and gender. In the study by Eisazadeh *et al.*, students from Qazvin were selected through cluster sampling. The results showed that the mean social anxiety score was 33.1 in girls and 32.6 in boys. A positive and significant relationship was also observed between mentalization and social anxiety, while a negative and significant relationship was found between mentalization and social anxiety as well. In fact, the results indicated that adolescents in social environments are not able to properly regulate their emotions and mental states, and as a result they develop social anxiety [21].

In contrast, the study by Aliakbari Dehkordi *et al.*, conducted among 277 students, found no relationship between gender and social anxiety scores, which is inconsistent with the results of the present study [22]. This discrepancy may be due to differences in the study populations. In the present study, patients with cleft lip and palate were examined, whereas the mentioned study evaluated general student populations.

According to the findings, the level of social anxiety among the patients was reported to be high. In the study by Hashemi Nosrat Abad *et al.*, a positive and significant relationship was found between body dysmorphic concerns and social anxiety, while negative and significant relationships were observed between body dysmorphia and self-esteem, as well as between self-esteem and social anxiety. Reduced self-esteem in individuals with body dysmorphic concerns may lead to the emergence or intensification of social anxiety, whereas increased self-esteem may reduce the severity of this disorder [23].

Similarly, in the study by Harouni *et al.*, university students from different academic levels under the age of 40 were examined. The findings showed a statistically significant

relationship between social anxiety and body image, such that students with lower body image scores had higher levels of social anxiety [24]. In another study by Mohammadi *et al.*, 226 university students from different academic levels were examined. According to the results, 71% of the students had social anxiety scores below 50, indicating no significant symptoms of social anxiety. Meanwhile, 6% of the students had scores above 80, placing them in the severe or very severe social anxiety category. A significant relationship was also observed between social anxiety and variables such as description, conscious action, non-judgment, non-reactivity, perfectionism, and orderliness [25].

In summary, previous studies indicate that negative body image and related psychological factors are associated with increased social anxiety. In line with these findings, the results of the present study also showed that the presence of cleft lip and palate—as a factor affecting physical appearance—may be associated with increased social anxiety.

Conclusion

The results of this study on 80 patients with cleft lip and palate showed that the mean level of social anxiety in this group was relatively high, with the highest score related to fear of negative evaluation. Social anxiety was also significantly higher among females, urban residents, and individuals who had not undergone reconstructive surgery. Therefore, demographic factors and treatment status may play an important role in the severity of social anxiety among these patients.

Reference

1. Tanhadoust K, Bezi M, Ghafarijam M, Amoojavadi F. The effect of group psychotherapy based on acceptance and commitment on loneliness and social adjustment in girls with social anxiety disorder. *J Nurs Educ.* 2021;9(1):32-42.
2. Baravordesh M, Chalabianloo G, Abdi R. The role of personality dysfunctions and abnormal personality traits in predicting symptoms of generalized anxiety. *J Psychol Stud.* 2025;21(2):39-55. doi: 10.22051/psy.2025.48899.3053.

3. Varaei S, Keshavarz S, Nikbakhtnasrabadi A, Shamsizadeh M, Kazemnejad A. The effect of orientation tour with angiography procedure on anxiety and satisfaction of patients undergoing coronary angiography. *J Nurs Educ.* 2013;1(2):1-10.
4. Shomali Ahmadabadi M, Zabihi M, Khodarahmi A, Barkhordari Ahmadabadi A. Mediating role of self-criticism in the relationship between fear of negative evaluation and self-compassion with social anxiety in Yazd medical students. *J Shahid Sadoughi Univ Med Sci.* 2024;32(8):8160-72. doi: 10.18502/ssu.v32i8.16856.
5. Tavangar Dehghan M, Sodagar S, Nayeibinia AS. The relationship between quality of life and pregnancy anxiety with the mediation of childbirth self-efficacy and social support. *J Ilam Univ Med Sci.* 2026;34(1):58-71.
6. Alikhani H, Samani S, Bardideh MR, Mozaffari M. Intervention based sociodrama: evaluating the effectiveness on the anxiety in children with social anxiety disorder. *J Nurs Educ.* 2022;10(3):88-103. doi: 10.22034/IJPN.10.3.88.
7. Sharifi P, Esazadegan A, Soleimani E. The structural model of the relationships between perceived stigma, fear of recurrence, and psychological flexibility with psychological distress in women with breast cancer. *J Psychol Stud.* 2025;21(2):57-72. doi: 10.22051/psy.2025.49337.3078.
8. Hoda J, Mohammadamin Khajeh Ahmad A, Zahra Dasht B. Cognitive rehabilitation improves memory and distress tolerance in social anxiety disorder: a quasi-experimental study. *Int J Health Stud.* 2025;11(2):28-34.
9. Fatemeh T, Hossein M, Abolfazl B. Comparison of the effectiveness of dialectical behavior therapy and Meichenbaum-based cognitive therapy on fear of negative evaluation in students with social anxiety disorder. *J Dev Neurol Disord.* 2025;4(1):111-7.
10. Rezazadeh Khalkhali F, Mehrinejad SA, Khodabakhsh Pirkalani R. The relationship between social anxiety disorder and attachment styles in students: the mediating role of social skills. *J Psychol Stud.* 2025;21(2):73-90. doi: 10.22051/psy.2025.49291.3074.
11. Hafez AA, Shahidi S, Nejat P, Moloodi R. The prediction model of social anxiety symptoms based on early maladaptive schemas with the sequential mediating role of cognitive fusion and experiential avoidance. *Rooyesh-e-Ravanshenasi J.* 2026;14(12):11-22.
12. Karimiazar F, Saken Azari R, Abdoltajedini P, Azmoudeh M. The effect of acceptance and commitment-based therapy on the symptoms of social anxiety in shy girls' adolescents. *J Nurs Educ.* 2022;10(1):46-54. doi: 10.22034/ijpn.10.1.46.
13. Farnoush F, Khodadadi M, Nouhi S. Comparing effectiveness of virtual reality therapy and mindfulness-acceptance therapy on attention focus and social anxiety symptoms. *Med Sci J Islam Azad Univ* 2020;30(3):299-312. doi: 10.29252/iau.30.3.299.
14. Ghaffari M, Narimani M, Basharpour S, Hajlo N. Comparison of the effects of personal intelligence and body intelligence training on the symptoms of social anxiety in cosmetic surgery applicants. *J Nurs Educ.* 2017;5(3):37-47. doi: 10.21859/ijpn-05036.
15. Hassanzadeh N, Azmoudeh M, Egbali A, Rezaei A. Body image concern in patients with psoriasis: comparison of two mindfulness-based stress reduction (MBSR) with and without transcranial direct-current stimulation (tDCS) treatment approaches. *J Nurs Educ.* 2025;13(3):94-107. doi: 10.22034/ijpn.13.3.94.
16. Usefi A, Taher M, Aghaei H, Bagherinia H. Body image anxiety in women; comparison of two cognitive-behavioral and interpersonal treatment approaches. *J Sabzevar Univ Med Sci.* 2023;29(6):833-46.
17. Ghobadi N, Karami S. A review of therapeutic approaches to cleft lip and cleft palate. *Clin Excel.* 2020;9(3):35-42.
18. Namdar P, Etezadi T, Raisolsadat SMA, Lal Alizadeh F, Shiva A. Treatment goals in individuals with orofacial clefts: a literature review. *Clin Excel.* 2020;9(4):45-55.
19. Homaei F, Khaleghipour S, Khoshakhlagh H. The effectiveness of psychodrama on emotional expression and social competence of adolescents with social anxiety. *Islam Life Style Cent Health.* 2022;6(3):255-63. Available from: <https://sid.ir/paper/1115490/en>.
20. Haller SP, Kadosh KC, Scerif G, Lau JY. Social anxiety disorder in adolescence: how developmental cognitive neuroscience findings may shape understanding and interventions for psychopathology. *Dev Cogn Neurosci.* 2015;13:11-20.
21. Eisazadeh M, Bahramkhani M. Predicting social anxiety based on attachment styles, mentalization capacity, and ego strength in adolescents of Qazvin city, Iran. *Iran J Psychiatry Clin Psychol.* 2024;30(1):949.1. doi: 10.32598/ijpcp.30.949.1.
22. Aliakbari Dehkordi M, Zare H, Asgharnejad Farid A, Hashemian R. Investigations of the relation between social anxiety, cognitive and behavioral aspect of it and sex and gender-role in adolescence. *Iran J Psychiatry Clin Psychol.* 2017;22(4):340-7. doi: 10.18869/nirp.ijpcp.22.4.340.
23. Mozafari A, Hashemi Nosrat Abad T, Mashinchi Abbasi N. Relationship between body dysmorphic disorder symptoms and social anxiety symptoms through self-esteem. *J Psychol.* 2026;29(4): [pages not provided].
24. Harouni B, KaH. The mediating role of self-compassion in the relationship between self-concept and body image and social anxiety in female students of Isfahan University of Technology. *Woman Interdiscip Res J.* 2022;4(1). doi: 10.1001.1.27831175.1401.4.1.4.0.
25. Mohammadi A, Roshan Chasli R. Structural relationships of social anxiety symptoms with perfectionism in students: the mediating role of mindfulness. *Clin Psychol Pers.* 2021;19(1):109-19. doi: 10.22070/cpap.2021.3301.

How to Cite This Article

Soltanmohamadi M. Assessment of social anxiety and associated factors in patients with cleft lip and palate in Ilam city in 2026. *International Journal of Medical and All Body Health Research.* 2026;7(2):67-70. doi:10.54660/IJMBHR.2026.7.2.67-70.

Creative Commons (CC) License

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0) License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.