



International Journal of Medical and All Body Health Research



International Journal of Medical and all body Health Research

ISSN: 2582-8940

Received: 01-01-2021; Accepted: 17-01-2021

www.allmedicaljournal.com

Volume 2; Issue 1; January-March 2021; Page No. 05-09

Prognostic Indicators in Patients with Intracerebral Hematoma in an urban clinical setting of a resource limited Country

Hugues Ghislain Atakla ^{1*}, Fatoumata Louncy Barry ², Mahugnon Maurel Ulrich Denis Noudohounsi ³, Benjamin Bekoe Ofosu ⁴, Sulaimi Sulemana Ummi ⁵, Dismand Stephan Houinato ⁶

^{1, 6}Department of Neurology, University Hospital Center Hubert Koutoukou MAGA, Cotonou, Benin

¹Laboratory of Noncommunicable and Neurologic Diseases Epidemiology, Faculty of Health Science, University of Abomey-Calavi, Cotonou, Benin

²Neurosurgery Department, University Hospital Center, Conakry, Guinea

³Monitoring and evaluation/WHO AFRO, Research Project Manager, Brazzaville, Congo

^{4, 6}Intensive Care Unit, MAB International Clinic, Accra, Ghana

Corresponding Author: **Hugues Ghislain Atakla**

Abstract

Context and Objective: Hemorrhagic Cerebrovascular Accident represents 10 to 15% of all strokes and are often related to the spontaneous rupture of small vessels weakened by chronic arterial hypertension, amyloid angiopathy, tumor bleed, arteriovenous malformation (AVM), and coagulopathies. The aim of this study was to determine probable prognostic indicators in functional outcomes of patient with ICH.

Patients and Method: This was a retrospective analytical study of all patients (men and women, of all ages) who were hospitalized at the Neurology Department of the University Hospital of Conakry for an intracerebral hematoma over the 24-month period. Only the records of patients in whom intracerebral hematoma was confirmed by brain imaging were included in this study. The variables studied were epidemiological, clinical, and paraclinical. Therapeutic, evolutionary and prognostic variables were also taken into account. Logistic regression (univariate and multivariate)

was used to identify the prognostic determinants of intracerebral hematomas at $p < 0.05$. Data were entered using Epi Info software version 7.1.4.0 and then analyzed using STATA/SE software version 11.2.

Results: We collected 105 cases of spontaneous cerebral hematomas confirmed by brain imaging, i.e. a frequency of 21%. Both sexes were found, with a male predominance of 60% and a sex ratio of 1.50. Arterial hypertension was identified as the cause in 94 patients (89.52%) out of the 105 patients analyzed in this series, followed by arteriovenous malformations in 6.67% of the patients, 2.86% of cases of amyloid angiopathy and 0.95% of unknown cause. During the hospitalization of the patients, we recorded 20% of deaths.

Conclusion: Previous quality of life, comorbidities affects the prognostic course in patient with ICH and should be taken into account in the prediction of disability and future quality of life of patients with intracerebral hematoma.

Keywords: Prognosis, Intracerebral hematomas, Limited health resources, Guinea

Introduction

Spontaneous cerebral hematoma is a specific type of stroke amongst the different groups of strokes that exist. They correspond to a bleeding in the cerebral parenchyma that can extend to the cerebral ventricles ^[1]. They represent 10 to 15% of all strokes ^[2] and are often related to the spontaneous rupture of small vessels weakened by chronic arterial hypertension or amyloid angiopathy ^[3]. More rarely, hemorrhage is secondary to rupture of a vascular malformation, tumor, venous thrombosis, hemorrhagic transformation of an arterial infarct, cavernoma, cerebral phlebitis, coagulopathy, or anticoagulant medication ^[3, 4]. Spontaneous cerebral hematomas are responsible for 30 to 50% of deaths and nearly 80% of patients are dependent at six months ^[3]. It is not related to gender and this frequency increases with age and half of the deaths occur within the first 48 hours ^[5]. The clinic is conditioned by the location of the hematoma. Deep hematomas (basal ganglia) are responsible for sensory and/or motor deficits in the hemisphere contralateral to the lesion ^[6]. While lobar bleeds are most often responsible for coma. Subtentorial hematomas are less frequent than supratentorial hematomas ^[6]. Close monitoring will focus on assessing the consequences of the oedema generated and look for signs of intracranial hypertension and for evidence of secondary hydrocephalus ^[7]. Diagnosis is made by brain imaging ^[5].

In the acute stage, the haematoma is typically hyperdense and the oedema peri lesional [4]. The treatment of haematomas aims to minimize the risk of further bleeding and the consequences of intracranial hypertension in order to reduce the risk of complications related to neurological symptoms [8]. The prognosis of ICH at six months is much less favourable because they cause severe neurological sequelae with only 20% of patients being independent at six months [9]. Recovery of motor disability is assessed by the Rankin score.

The Rankin handicap scale was developed on the basis of the prognosis of patients who have had a stroke [9, 10]. Certain clinical and radiological indicators, including Hemphill's "ICH Score" or cerebro-meningeal hemorrhage score, predict mortality at 30 days and functional prognosis at one year. A modified ICH Score of less than 2 is an excellent predictor of good prognosis [5]. The NIHSS (Stroke Severity Scale) score is a better predictor of morbidity [5, 11]. Thus, this series allows us to evaluate the prognosis of intracerebral hematomas at the neurology department of Ignace Deen University Hospital Center in Conakry.

Patients and Method

This was an analytical retrospective study, conducted on all patients who were hospitalized for an intracerebral hematoma over a period from January 1, 2017 to December 31, 2019. All records of patients hospitalized with intracerebral hematoma confirmed by brain CT and/or brain MRI were included in this study. We did not include records of patients who had intracerebral hematoma but refused to participate in the study, and patients who we could not reach to reassess the prognosis. The following data were collected: age; sex; socio-professional strata, marital status; origin; history; risk factors; secondary systemic brain injury (SSBA); mode of onset; time to consultation; etiologies; initial assessment of stroke severity; complications and paraclinics; therapeutic variables; evolutionary variables; prognosis.

Initial assessment of the severity of intracerebral hematoma

The assessment of severity was done mainly by a clinical scale, the NIHSS (National Institute Health Stroke Scale) score, and by prognostic factors mainly studied in the situation of severe stroke. The NIHSS scale is rated from 0 (normal) to 42 (maximum score). It is composed of 11 items exploring consciousness, oculomotricity, visual field, motor skills, sensitivity, ataxia, language and neglect [6]. A stroke is defined as:

- Minor if the score is between [0 and 5]
- Moderate for a score between [6 and 10].
- Severe if the score is between [11 and 15].
- Very severe if the score is between [16 and 20].
- Severe for a score higher than 20.

The evolutionary variables (favorable and death) were defined by the following criteria:

- NIHSS score at admission: was used to assess the severity of acute stroke and a fairly reliable predictor of residual disability.
- Hematoma volume: was calculated according to the formula $A \times B \times C / 0.5$ and expressed in milliliters (A is the largest diameter of the hematoma, B is the largest diameter perpendicular to A on the same slice, C is the product of the number of slices in which the hematoma

is seen by the thickness of the slices). The average size of the hematoma was determined

- ICH scores at admission: performed at admission has a predictive value for one-month mortality:

Table 1: HEMPHILL'S SCORE (ICH)

Hemphill's Score (ICH)		
Criteria		Scores
Glasgow coma scale	3-4	2
	5-12	1
	13-15	0
Volume of the hematoma	≥ 30 ml	1
	< 30 ml	0
Ventricular effusion rate	Oui	1
	Non	0
Age (years)	≥ 80 ans	1
	< 80 ans	0
Infratentorial topography	Yes	1
	No	0
Mortality at 30 days		
	≥ 5	100%
	4	97%
	3	72%
	2	26%
	1	13%
	0	0%

- RANKIN admission score: this was performed on all our patients. It is a global evaluation scale of handicap, performed in 5 minutes. This score allows us to classify hemiplegic patients into six categories according to their ability to walk and their dependence on activities of daily living:

- 0: No symptoms
- 1: No disability apart from symptoms: activities and autonomy maintained
- 2: Low disability: inability to perform usual activities but autonomy;
- 3: Moderate disability: needs help but can walk without assistance
- 4: Moderately severe disability: walking and daily activities impossible without assistance.
- 5: Major disability: permanent bed rest, incontinence and permanent nursing care
- 6: Death

The information collected in this study was obtained with the consent of the patients and/or their parents. The anonymity and confidentiality of the information collected were preserved. All the data were entered using Epi Info software version 7.1.4.0 and then analyzed using STATA/SE software version 11.2.

Results

During the study period, we collected 105 cases of spontaneous cerebral hematomas confirmed by brain CT and brain MRI, out of a total of 491 stroke patients, representing an incidence of 21%.

Diabetes is a global health problem and is a common endocrine problem associated with multiple disorders that include metabolic, cellular and blood disturbances. Type-II diabetes is a multi causal disease which develops slowly and initially, it commences with insulin resistance, which progress gradually with the passage of time




Fig 1: Hospital incidence of spontaneous cerebral hematoma.

Male and female gender were found with a male predominance of 60% and a sex ratio of 1.50. The mean age of our patients was 56 ± 15 years with extremes of 22 and 93 years. The age group most affected by the disease was that between 60 and 70 years with 25 patients or 23.81%. We noted a clear predominance of married patients with 96%, followed by widows with 6.67%, singles and widowers respectively 0.95%. Seventy-three (73%) of the patients were from urban areas, compared to 27% from rural areas. Of the 105 patients analyzed in this series, 89 (84.76%) had a comorbidity of arterial hypertension, 10 (9.52%) had type II diabetes, 5 (4.76%) had a previous ischemic stroke and only 1 (0.96) patient had heart disease.

Table 2: Distribution of patient records by history

History	Staff	Percentage
High BP	89	84.76
Type 2 diabetes	10	9.52
Cardiopathy	1	0.96
History of Ischemic Stroke	5	4.76
Total	105	10

At admission, 39 patients (37.14%) had an NIHSS score of 11-15; 36 (34.29%) had an NIHSS score of 16-20. Only 5 or 4.76% of the patients had an NIHSS score of 0 - 5, i.e. a mean NIHSS score of 14.18 ± 4.72. All patients had a brain CT scan and among them, 30 patients (28.57%) also had a brain MRI scan. The deep location was found in sixty-five (65) or 61.9% of patients, 13 or 12.38% lobar, 12 or 11.43% in the brainstem and intraventricular in 15 or 14.29% of cases with associated lesions in 17 cases or 15%. We found associated lesions in 17 patients, including 12 cases of involvement (11.43%), 6 cases of mass effect (5.71%) and 5 cases of ventricular dilatation (4.76%).

Table 3: Distribution of patient charts according to lesion topography on imaging

Topography of the injury	Staff	Percentage (%)
Deep	65	61.9
Lobar	13	12.38
Brainstem	12	11.43
Intraventricular	15	14.29
*Associated lesions	17	15

* Mass effect: 6; *Hydrocephalus: 5; Engagement: 12

The mean volume of the hematoma was 43.19 ml ± 35.047.

Table 4: Distribution of patient charts by CT hematoma size

Volume of hematoma on CT scan	Staff	Percentage (%)
≤30ml	73	69.52
>30ml	32	30.48
Total	105	100

The functional prognosis of our patients were assessed by the Rankin score. 81 patients equivalent to 77.14% had a Rankin score of 4 on admission, followed by 13 or 12.38% with a score of 3; only 1 patient or 0.95% had a score of 6, with a mean Rankin score of 3.85 ± 0.68 on admission.

47 patients or 44.76% had an admission ICH score of 2; 25 patients or 23.81% had an ICH score of 1 and 3 respectively; 2 or 1.90% had a score of 4 and only 1 patient had a score of 5, the mean admission ICH score was 1.97 ± 0.91. Hyperthermia and hyperglycemia were the secondary brain attacks of systemic origin occurring during hospitalization, distributed respectively in the following proportions: 39 patients (37.14%), 24 patients (22.86%) (Table 5). The complications occurred were dominated by inhalation pneumonitis and pressure sores, represented respectively as follows: 15 (14.28%), 5 (4.76%). (Table 6)

The evolution was favorable in 84 patients or 80% in contrast to 21 patients or 20% death (Figure 2). Length of stay ≥10 days was associated with high mortality with a p value = 0.001. Hematoma volume was associated with high mortality with a p value = 0.02. On the other hand, GCS score <7 on admission was correlated with high mortality with a significance value = 0.001. In this study the high ICH score was associated with high mortality, we found a significance threshold with a p value = 0.0001.

Table 5: Distribution of patient folder according to secondary cerebral attack of systemic origin.

SCASO	Staff	Percentage (%)
Hypoglycemia	5	4.76
Hyperthermia	39	37.14
Hyperglycemia	24	22.86

Table 6: Distribution of patient folder according to complications during hospitalization.

Complications	Staff	Percentage (%)
Pressure ulcer	5	4.76
Inhalation pneumopathies	15	14.28
Septic shock	3	2.85
Recurrence	0	0
Pulmonary embolism	2	1.90




Fig 2: Distribution of patient folder according to progression

Discussion

Our study found an overall frequency of 21%. Our results are superior to those of Basse A et al in 2017² who found 10 to 15% of strokes.

Both sexes were found with a male predominance of 60% and a sex ratio of 1.50. Our result is lower than that of COLNAT-COULBOIS S. et al. at Nancy University Hospital in France, who reported a clear male predominance of 62%¹². This result could be explained in our context by the fact that the male sex is most exposed to cardiovascular risk factors before the age of 55 or less. The mean age of our patients was 15.12 years with extremes of 22 and 93 years. The most affected age group was between 60 and 70 years old. Our results are consistent with those of Bilongo Manene¹³ who found a mean age of 61 years. These results show that the incidence of hemorrhagic stroke increases with age. We noted a clear predominance in married patients with 96% followed by widows with 6.67%, singles 0.95%. This result is comparable to that of Sene Diouf. F et al. at Fann University Hospital in Dakar¹⁴ who reported 35 (68.5%) married cases followed by 9 (17.6%) widowed cases. This could be explained by the fact that married people are more exposed to family stress than unmarried people. At admission 39 patients or 37.14% had an NIHSS score between 11-15; 36 patients or 34.29% had an NIHSS score of 16 - 20. Only 5 or 4.76% of the patients had an NIHSS score between 0 and 5 with a mean NIHSS score of 14.18 ± 4.72 . These results are higher than those of CISSE F.A et al¹⁵ who found a mean NIHSS score of 10.6 ± 4.3 . Deep localization was found in 65 patients (61.9%), lobar in 13 (12.38%), brain stem in 12 (11.43%) and intraventricular in 14.29%. Associated lesions were found in 17 patients, including 12 cases of engagement (11.43%), mass effect (5.71%) (mass effect 6) and 5 cases of ventricular dilatation (4.76%). Our results are similar to those of Kanté M¹⁶ which found 48.6% deep localization, 29.2% lobar localization, 9.7% brain stem and intraventricular localization in 12.9%, with associated lesions in 15.3%. The mean volume of the hematoma was $43.19 \text{ ml} \pm 35.047$. This result is close to that of Lelo Tshikwela et al¹⁷ who found a mean volume of the hematoma of 37 ml. Hypertension was the cause found in 89.52% of patients; followed by arteriovenous malformations in 6.67% of patients, 2.86% amyloid angiopathy and 0.95% unknown cause. Our result is superior to that of Jayantee Kalita et al in India¹⁸ who reported in their study 79.2% of cases of hypertension as the first cause.

All patients followed physiotherapy sessions for the rehabilitation of hemiplegia. Forty (40) patients, i.e. 38.09%, underwent medicalized rehabilitation; 35, i.e. 33.33% of the patients underwent mixed rehabilitation and 30, i.e. 28.57% underwent non-medicalized rehabilitation. Our result is different from that of Sene Diouf et al. in Dakar¹⁴ who reported in their study 53.5% of cases of rehabilitation carried out by qualified personnel; 29.4% by unqualified personnel and 24.1% by a family member. This result could be explained by the fact that unlike developed countries where the treatment of hemiplegia of vascular origin is global and multidisciplinary, in developing countries it is limited to physiotherapy sessions. Occupational therapy is almost non-existent and speech therapy rehabilitation is rare and only exists in private practice Sene Diouf F et al in Dakar¹⁴.

Evaluation of the functional prognosis of our patients was assessed by the Rankin score. 81 patients or 77.14% had a Rankin score of 4 at admission, followed by 13 or 12.38%

had a score of 3; only 1 patient or 0.95% had a score of 6, with a mean Rankin score at admission of 3.85 ± 0.68 . On admission, 44.76% of patients had an ICH score of 2; 25 patients or 23.81% had an ICH score of 1 and 3 respectively; 2 or 1.90% had a score of 4 and only 1 patient had a score of 5. The mean HCI score at admission was 1.97 ± 0.91 . Our result is superior to that of A. Gnazebo et al¹⁹ who found an ICH score that varied between (0 and 4) with a mean ICH score of 1. The volume of the haematoma was associated with high mortality with a p value = 0.02. This result is comparable to that of Capellari et al. in 2015²⁰ who reported a hematoma volume \geq of 30 ml with a significance threshold < 0.001 .

The Glasgow score < 7 at admission was correlated with high mortality with a significance value $p = 0.001$. Rasamoelina N et al in 2018 in their study reported that the Glasgow score ≤ 8 was correlated with high mortality with a significance value of $p = 0.03$ ²¹. Our result could be explained by the fact that the management of intracerebral hematomas in the neurovascular unit during the acute phase would considerably reduce their mortality. In this work, the high ICH score was associated with high mortality, and we found a threshold of significance with a value of $p = 0.0001$. This result could be explained by the fact that the components of the ICH score are predictive factors associated with high mortality during ICH. The evolution was favourable in 84 patients, i.e. 80% as opposed to 21 patients, i.e. 20% of deaths. This high mortality rate could be explained by the fact that our deceased patients had an ICH score between (3 and 5) on admission. This result could be explained by the fact that an extended hospital stay increases the risk of medical complications.

Many studies describe the functional prognosis and early complications of intracerebral hematomas. However, very little work has been done in this area in developing countries such as ours. We believe that a future study on the follow-up of the CT evolution of intracerebral hematomas will facilitate a better understanding of the matter and guide the clinician in therapeutic decisions.

Conclusion

Spontaneous brain hematomas usually have a more severe prognosis; they also cause severe neurological sequelae. Morbidity and mortality are largely influenced by complications in the acute phase. Age above 80 years; NIHSS score above 15 at admission; and the occurrence of pressure ulcers and inhalation pneumopathies were factors in poor prognosis. Education of the population to consult at the onset of the first signs is necessary. The organization of care channels and an early start of functional rehabilitation are key elements of good prognosis. It is therefore important to take this into account when assessing the neurological prognosis. Finally, previous autonomy and comorbidities also modify the prognosis and should be taken into account in the prediction of disability and future quality of life.

AVM: arteriovenous malformation

ICH: Intracerebral hemorrhage

ICH score: Intracerebral hemorrhage score

NIHSS score: National Institutes of health stroke Scale/score

SCASO: Secondary cerebral attack of systemic origin

Acknowledgement

The authors would like to thank the patients and the ethics committee for consenting to this study.

Authors' contributions:

All authors contributed to this work.

Availability of data and materials

All data sets used and/or analyzed in this study are available from the corresponding author upon reasonable request.

Funding

The authors did not receive any funded support for research, writing and publication of this article.

Declaration of interest

All the authors do not have any possible conflicts of interest.

References

1. Qureshi A, Tuhim S, Broderick JP, *et al.* Spontaneous intracerebral hemorrhage. *N Engl J Med.* 2001; 344(19):1450-60. doi: 10.1056/NEJM200105103441907.
2. Basse AM, Sow AD, Toure K, *et al.* Épidémiologie de l'accident vasculaire cérébral hémorragique chez les patients hospitalisés à la clinique neurologique du CHU de Fann (Dakar). *Revue neurologique.* 173 S(2017) S81–S132
3. Crozier S, Traitement des hémorragies Cérébrales. Service urgences cérébrovasculaires. Groupe hospitalier Pitié-Salpêtrière. Paris (75) tome 36- n°1 jan 2007.
4. Hamon M, Leclerc X, Oppenheim C, Gauvrit Jy., Meder Jf, Pruvo JP. Sémiologie TDM et IRM des hématomas intracérébraux: évolution chronologique. *Rev Neurol (Paris)* 2005, 161:10,997-1006. Doi: RN-10-2005-161-10-0035-3787-101019-200505313
5. Hermitte Y, Do-Esposito S, Foudi H, Tazarourte K. Accident vasculaire cérébral hémorragique. 53 congrès national d'anesthésie et de réanimation. Infirmiers. Infirmier (e)s d'urgence, 2011. Sfm
6. Chung CS, Caplan LR, Han W, *et al.* Thalamic hemorrhage. *Brain.* 1996; 119(Pt 6):1873-86. doi: 10.1093/brain/119.6.1873
7. Kumral E, Kocaer T. Thalamic hemorrhage A prospective study of 100 patients. *Stroke.* 1995; 26(6):964-70. doi: 10.1161/01.str.26.6.964
8. Mayer SA, Rincon F. Treatment of intracerebral hemorrhage. *Lancet Neurol.* 2005; 4(10):662-72. doi: 10.1016/S1474-4422(05)70195-2.
9. Agency for Health Care Policy Research. Post-stroke rehabilitation: assessment, referral, and patient management. Quick reference guide for clinicians. Rock-ville (MD): AHCP; 1995.
10. Duncan P, Lai S, Van Culin V, *et al.* Development of a comprehensive assessment toolbox for stroke. *Clin Geriatr Med.* 1999;15(4):885-915.
11. Teitelbaum J, Scarborough F, et François S. *Le Médecin du Québec.* 2006; 41:6.
12. Morgensen LB, Anderson G, Hemphill, *et al.* Guidelines for the management of spontaneous Intracerebral. *Stroke* 2010; 41:2108-2129. <https://doi.org/10.1161/STR.0b013e3181ec611b>
13. Bilongo-Manene B. Mortalité et morbidité des AVC dans le service d'anesthésie réanimation du CHU Gabriel Touré. These, Med, Bamako, 2007, n°137.
14. Sene Diouf F, Mapoure NY, Ndiaye M, *et al.* Pronostic des hémorragies intracérébrales avec coma dans une unité de neuroréanimation tropicale. *Med Trop* 2008; 68:606-610.
15. Cisse FA, Balde AT, Sakadi Foksouna et al. Hématomes intracérébraux: 129 cas observés au service de neurologie du CHU de Conakry. *revue neurologique* 174 2018, S55-S108
16. Kanté M. Pronostic des hématomas intracérébraux spontanés au service de neurologie de l'hôpital Ignace Deen. thèse de doctorat en médecine service de neurologie CHU Conakry, 2014.
17. Lelo Tshikwela M, Longo-Mbenza B. Accident vasculaire cérébral hémorragique du noir africain: caractéristiques scanographiques des hématomas intracérébraux à Kinshasa. *J Afr Imag Méd.* 2011; (4)7:355-362. DOI: 10.11604/pamj.2015.21.160.6852
18. Jayantee Kalita, Gourav, Pankaj K, et al. intracerebral hemorrhage in young from a tertiary neurology center in North India; *Journal of the Neurological Sciences* 336 (2014) 42–47. DOI:<https://doi.org/10.1016/j.jns.2013.09.037>
19. Gnazebo A, Kouame-Assouan AE, Akani AF et al ; Score ICH et pronostic vital des hématomas cérébraux au service de neurologie du CHU de Bouaké. *revue neurologique.* 2018; 174:S55-S108. Doi: 10.1016/j.neurol.2018.01.181
20. Cappellari M, Zivelonghi C, Moretto G, *et al.* The etiologic subtype of intracerebral hemorrhage may influence the risk of significant hematoma expansion; *Journal of the Neurological Sciences* 2015; 359:293-297. DOI: <https://doi.org/10.1016/j.jns.2015.11.024>
21. Rasamoelina N, Andrianimaro FM, Randriamarolahy A. Predictive factors of mortality of hemorrhagic stroke observed in the intensive care unit of the teaching hospital tambohobe, fianarantsoa. *Paripex - Indian Journal of Research.* 2018; 7(1). | Print ISSN No 2250-1991