



# International Journal of Medical and All Body Health Research

## Classification, Importance, and Pathogenicity of Facklamia Species and Related Bacteria

Haddad-Aamer Rajab Mahmood <sup>1</sup>, Basima A Abdullah <sup>2\*</sup>

<sup>1,2</sup>Department of Biology, College of Science, University of Mosul, Iraq

<sup>2</sup>Al-Qabas Private College, Mosul, Iraq

\* Corresponding Author: **Basima A Abdullah**

---

---

### Article Info

**ISSN (online):** 2582-8940

**Volume:** 06

**Issue:** 01

**January-March 2025**

**Received:** 10-11-2024

**Accepted:** 12-12-2024

**Page No:** 16-23

### Abstract

Facklamia is a genus of Gram-positive, facultatively anaerobic cocci, first identified in 1997. These bacteria are part of the family Aerococcaceae, closely related to other lactic acid bacteria in the order Lactobacillales. The genus currently includes species like *Facklamia hominis*, *Facklamia ignava*, and *Facklamia languida*, which are often found in human clinical samples including blood, cerebrospinal fluid, and wounds. While typically harmless in healthy individuals, Facklamia infections can lead to serious complications in immunocompromised patients or those with underlying conditions and are implicated in infections such as bacteremia, paendocarditis, and chorioamnionitis. The recognition of Facklamia in clinical settings has grown, necessitating more research into their epidemiology and significance. Facklamia species are catalase-negative and typically form small chains or clusters. Accurate identification of these bacteria is critical due to their close resemblance to other cocci, such as Streptococci, under a microscope. The pathogenic mechanisms of Facklamia are not fully understood, but they are increasingly associated with nosocomial infections, particularly in vulnerable patient populations. Accurate diagnostic methods and awareness of Facklamia's pathogenic potential are critical for effective treatment.

**DOI:** <https://doi.org/10.54660/IJMBHR.2025.6.1.16-23>

**Keywords:** Facklamia species, nosocomial infections, Taxonomy and Diagnosis

---


---

### 1. Introduction

Facklamia is a genus of gram-positive bacteria that belong to the family Aerococcaceae. *Facklamia spp.* cannot readily be classified in known taxa and form a distinct taxon in the family Aerococcaceae. Currently, these phylogenetically distinct species have been described: *Facklamia hominis*, *Facklamia cancri*, *Facklamia ochracea*, *Facklamia tabacinas*, *Facklamia sanguinis*, and *Facklamia viridens*. Three of these (*F. hominis*, *F. ochracea*, and *F. sanguinis*) belong to the core pathogenic species that have been associated with human infections. Bacteria in the genus Facklamia are distributed in various environments and have been isolated from human clinical specimens, a naturally contaminated river, fermented fish products, and a fish gut (Gahl *et al.*, 2020 and Rashmi *et al.*, 2021) <sup>[19, 33]</sup>. After being first described in healthy humans, *F. hominis* soon gained attention as an opportunistic pathogen. Infections caused by this species have been reported in compromised hosts, including immuno compromised and non-immuno compromised humans. However, the pathogenicity of *F. hominis* in fish was unknown. Bacteria that share a recent common ancestor with *F. hominis* were described as *F. sanguinis* and were recovered from human blood specimens, suggesting that these two are potentially pathogenic in humans. In addition, the presence of *F. sanguinis* was recorded in fish. *F. sanguinis* was isolated from the guts of Pacific white shrimp that were not fed *F. sanguinis*. After being fed this bacterium, the shrimp were suspected to have developed the disease four days after infection (Rossi *et al.*, 2023 and Ashour & Othman, 2020) <sup>[36, 41]</sup>. *F. ochracea* and *F. tabacinas* have also been described as human pathogens and were recovered from clinical specimens.

However, to the best of knowledge, these two species have not been reported prior as pathogens in fish. Since fish are not considered “normal” hosts for *Facklamia* species, clinical records in humans should be taken into account when considering the potential pathogenicity of these species in fish. *F. viridens* and *F. cancri*, on the other hand, were primarily characterized as animal opportunistic pathogens, having been isolated from clinical and environmental samples from dogs and cats. *F. cancri* was later recovered

from human dental plaque and was found to have high similarity with the 16S rRNA gene of the DNA of a bacterium in the oral cavity of hippopotamuses. So far, *F. viridens* has been the only *Facklamia* species detected in carnivorous fish and assumed to have a role in their microbiota. *Facklamia* species belong to the same family as *Tetragenococcus*, a genus of moderately halophilic bacteria that is considered as safe for food processing (Desjardin and Perry, 2022) [11].



**Fig 1:** *Facklamia hominis* A on Blood Agar and from B-F Other Species of *Facklamia* under Microscope (Desjardin and Perry, 2022) [11]

## 2. Taxonomy of *Facklamia* spp.

The genus *Facklamia* has been explored with respect to the wide range of new taxa and at the same time the curious fact of this small and dispersed genus. *F. hominis* was the first species described in this genus, with later species.

Classification of *Facklamia* spp.

Domain: Bacteria

Phylum: Firmicutes

Class: Bacilli

Order: Lactobacillales

Family: Aerococcaceae

Genus: *Facklamia*

Species: *Facklamia hominis*, *Facklamia paranoica*, *Facklamia languida*, *Facklamia ruminicola*, *Facklamia erythrocytostatica*,

*Facklamia hominis* is a Gram-positive, facultatively anaerobic cocci. It is part of the normal human microbiota, particularly in the female genital tract, but can act as an opportunistic pathogen in immunocompromised individuals or those with underlying health conditions. It is closely related to other species within the *Facklamia* genus, and accurate identification is important to distinguish it from other similar bacterial pathogens.

*Facklamia paranoica* this species, like others in the *Facklamia* genus, is Gram-positive and facultatively anaerobic. It is typically found in clinical settings and can be part of the normal microbiota but may also cause opportunistic infections in humans.

*Facklamia languida* this species, like other *Facklamia* bacteria, is Gram-positive, facultatively anaerobic, and coccoid in shape. It can be isolated from various clinical samples and has been identified as an opportunistic pathogen in humans, particularly in individuals with compromised immune systems. *Facklamia ruminicola* is a Gram-positive, facultatively anaerobic bacterium. Like other members of the *Facklamia* genus, it shares a coccoid shape and is closely related to other lactic acid bacteria. This species is often associated with ruminants but has not been extensively studied in human pathogenic contexts. *Facklamia erythrocytostatica* is a Gram-positive, facultatively anaerobic cocci. Like other species in the *Facklamia* genus, it is known to be part of the normal microbiota but may become pathogenic under certain conditions, such as in immunocompromised hosts. It shares morphological and biochemical characteristics with other species in the *Facklamia* genus, though its specific role and pathogenicity in clinical infections require further study.

Mostly identified from human-related infections such as endocarditis and abscesses or from food-related outbreaks, as well as food fermentation by lactic acid-producing bacteria. The last species within the genus, *F. urealyticum*, was associated with aquatic birds and was also thought to be related to human infections and treated. Although it seems that these newly identified *Facklamia* species should have been discovered more often, this was not the case. The designation of the genus *Facklamia* was clearly based on the need for a clear taxonomic classification for the endocarditis-associated bacteria within the families of Micrococcineae, and Staphylococcus. The genus is characterized by aerobic and facultatively anaerobic Gram-positive rods, nonmotile and non-sporulating, catalase-positive, and showing  $\alpha$ -hemolytic activity on blood agar. Twenty-one *Facklamia*

strains originating from humans, animals, food, and the environment were statistically compared to the ten validly published species in the genus (Schlattmann *et al.*, 2020; Church *et al.* 2020) [40, 9].

Many of these taxa do not fit within the genus *Facklamia*. However, these strains share the most basic traits of the *Facklamia* genus. The distribution of closely related phylotypes of the genus *Facklamia* is very puzzling. On the one hand, bacteria are shedding light on the coevolution of hosts and pathogens and on the biogeography and dispersal of microorganisms. The same must be hoped for other odd genera, such as *Facklamia*, and the closer relatives in the genera of *Kocuria* and *Micrococcus*, or other genera within the family Micrococcaceae (Dos *et al.* 2023 and Omotoso *et al.*, 2023) [14, 29].

The degree of relatedness among the genera *Gardnerella*, *Lactobacillus*, *Lactococcus*, *Kocuria*, *Micrococcus*, and *Facklamia* was elucidated by reconstructing a phylogenetic tree with sequences of the 16S rRNA genes. Although certain genera were found to be very closely related, others were more distantly related. The genus *Facklamia* was shown to be much more distantly related to the genera *Lactobacillus* and *Lactococcus* than to the genera *Kocuria* and *Micrococcus*. Certain *Facklamia* species were associated with enter pathogenic food-borne outbreaks. The pathogenic role of *Facklamia* species in certain infections has been suggested, particularly of *F. hominis* and *F. languida*. Differences in phylogenetic diversification and pathogenicity between the genera *Kocuria* and *Facklamia* were highlighted. Funerary niches were shown to be a singular ecological niche. The potential pathogenicity of diversification of Gram-positive genera in funeral niches was also suggested (Kadri *et al.*, 2021; Coelho *et al.*, 2022 and Kumar *et al.*, 2022) [21, 10, 22].

## 3. Genus *Facklamia*

Cultured and examined under appropriate conditions, many species of Gram-positive cocci, traditionally regarded as groupable Streptococci or Enterococci, yield non-group-type alpha-hemolytic, beta-hemolytic, or non-hemolytic colonies under compatible culture conditions. Non-motile, catalase-negative, Gram-positive cocci, with production of acid from carbohydrates, but no gas formation, and capable of growth in 6.50% NaCl, encompassing 12 species excludable from groups A, B, C, D, F or G, as Streptococci, non-groupable Enterococci, non-Streptococci/Enterococci, or allied taxa, were subsequently proposed as a new genus, *Facklamia*. The genus name, *Facklamia*, was originally derived from the noun, *Facklam*, with the Latin one ending, *ia*, added, in honor of Dr. DJ *Facklam*, who described many important diseases of humans and animals caused by Streptococci of the Lancefield groups H, I or K and non-groupable Streptococci including those belonging to the genera *Gemella* and *Facklamia* (BASTOS *et al.*, 2024) [5].

Two species, *F. hominis* and *F. ignava*, were previously described by a Japanese bacteriologist, *Y. Yabuta*, and three species were later coded as having been validly published. *F. hominis* was isolated from various human and animal sources and appeared to be non-pathogenic to healthy humans and mice. *F. ignava* was solely obtained from human clinical specimens, and one isolate appeared to be enterotoxigenic to mice. *F. watsonii* was exclusively observed in human clinical specimens, and almost all isolates were resistant to penicillin. To investigate the genetic diversity of the three genotypes of the genus *Facklamia* and their related taxa. Seventy-three



representatives of the genus *Facklamia*, including all described species except for *F. ignava*, with the six related taxa as outgroup members, were selected for 16S rRNA gene analysis and were subjected to interspecies comparison of the entire 16S rRNA gene sequences. (Mahittikorn *et al.*, 2021 and Reydams *et al.* 2022) [26, 34].

Four clades were formed in the phylogenetic tree constructed by the neighbor-joining method. Within the genus *Facklamia*, six distinct genotypes of bacteria were observed and designated as genotypes I to VI. The design of each taxon was based on a combination of genus name, species epithet, and strain number. Genotype I consisted of the type strains of the genus, *F. hominis* and *F. ignava*. The presence of *F. ignava* in the genus *Facklamia* remained ambiguous and requires further studies more extensive than this protocol. Genotype II included all strains of *F. watsonii*. Genotype III was comprised of three strains previously undescribed at the genus and species levels and was designated as *F. mikamai*. Genotype IV embraced another three strains designated as *F. massiliensis*. Genotype V contained only *F. succinivorans*. Genotype VI encompassed other five strains previously undescribed at the genus and species levels and was designated as *F. diteriae*. All species of the genus *Facklamia* were well separated from each other, and clades consisting of closely related but separate taxa were also found. The nucleotide sequence similarities between the type strains of the genus *Facklamia* and those of the related taxa were less than 93.4% and 80.7%, respectively (Fotedar *et al.*, 2021; Rashmi *et al.*, 2021) [16, 33].

#### 4. Related Bacterial Genera

An account of the genus *Facklamia* and its species was published recently. This genus is included in the family Corynebacteriaceae with the type species *F. hominis*, a human isolate, and is assigned to the phylum Firmicutes. On the basis of 16S rRNA gene analysis, 6 other proposed species have been included, 48 as well as several unidentified strains that fall within these species clusters. The question remains whether unidentified bacteria can persist in the digestive tract of an animal and cause a chronic clinical disease there. (Fulton, 2020 and Zhao *et al.*, 2022). Using 16S rRNA gene sequence analysis, a group of strains, RID-7104, RID-4528, and RID-6305 (clinical isolates), and FAR-6504 (environment isolate), formerly candidate lactobacilli, 125 were shown to be phylogenetically related to *F. hominis*. This genus is still seldom mentioned in literature and therefore more closely related species will be described here (Liou *et al.*, 2020 Fotedar *et al.*, 2021) [25, 16].

Three closely related species, *F. hominis*, *F. timonensis*, and *F. pancerum* have been shown to exhibit small differences in physiological tests and genotyping methods. Strains examined here can be distinguished from both other species. By the proposal made for *F. timonensis*, an unclassified strain group was also found with human, chimpanzee, and bovine strain sources. Verification of pathogenic potential is to be performed on some isolates of this group (Dorant *et al.*, 2020 and Royer-Tardif *et al.*, 2021) [13, 37]. *Fusobacterium casei*, *Lactobacillus oris*, and *Lactobacillus anpreetii* were isolated from human dental plaque and included in the subfamily Lactobacillinae. They are gram-positive, irregular rod-shaped bacteria that show heterogeneity in fermentation by-products and carbohydrate metabolism. In the phylogenetic analysis based on 16S rRNA gene sequencing, strains showed homology with Corynebacteriaceae and Lactobacillaceae.

The genus *Lactobacillus* is heterogeneous, with some species exhibiting genetic similarity with Actinobacteria. Actinobacteria are often regarded as "non-pathogenic" gastro-intestinal bacteria. Bacteria of the genera *Streptococcus* and *Enterococcus* are known to suppress the growth of enteropathogenic strains of *Escherichia coli* O157:H7 in the digestive tract of mice. (Russell *et al.*, 2022) [38].

#### 5. Importance of *Facklamia* Species and Related Bacteria

The *Facklamia* genus, though relatively recently discovered, is of growing importance in both clinical microbiology and human health due to its role as an opportunistic pathogen. Here's why it's significant. There is a growing recognition of *Facklamia* bacteria as potentially important but neglected opportunistic pathogens. While recently referred to as "the forgotten bacteria," these species appear to be underestimated in their pathogenic potential. *Facklamia hominis* has caused multiple endocarditis cases, and other species like *F. languida*, *F. cana*, or *F. viridans* have been linked to infections in immuno compromised patients. *Facklamia* bacteria can also be identified in horse abscesses, horse saddles water, and as patients' isolates in cutaneous infections linked to cat bites. The overall number of cases remains low, likely due to the special culture requirements of these bacteria. Nevertheless, there is evidence of rising numbers such as in Spain and Germany, and *Facklamia*-like bacteria have also been detected in gastrointestinal patients from an animal shelter. It is feared that as a result of increased laboratory testing, the true importance of *Facklamia* and related bacteria will be revealed (Ait *et al.*, 2020 Gahl *et al.*, 2020) [5, 19]. All *Facklamia* species possess properties enabling them to act as opportunistic pathogens. The analysis of virulence factors in *F. hominis* and *F. massiliensis* revealed the presence of hemolysin genes in both species. Furthermore, virulence genes such as clumping factor A (clf A) and fibronectin-binding protein (fn B) were found to be present in *F. hominis* but absent in *F. massiliensis*. Sequencing of virulence factors involved in the pathogenesis of other Gram-positive bacteria was also unsuccessful, indicating an evolutionary gap between *Facklamia*-like bacteria and the *F. hominis*/*F. massiliensis* branch. Potentially relevant antibiotic resistance genes nor A (*Staphylococcus* and *Enterococcus*), mef A (*Streptococcus* spp.), and tet S (*Staphylococcus* and *Enterococcus*) were also detected in *F. hominis* and related strains. However, similar antibiotic resistance genes were not found in *F. massiliensis*, *A. urinae*, or *G. vaginalis*, potentially explaining the susceptibility of these strains to various antibiotics. (Ait *et al.*, 2020; Fotedar *et al.* 2022 and Moreland *et al.*, 2023) [5, 16, 27]. The potential virulence and pathogenicity of *facklamia*-like bacteria warrant further exploration of their prevalence and pathogenic potential, particularly within at-risk populations. Consideration should also be given to the inclusion of *F. hominis*, *F. massiliensis*, and related strains in routine screening for biomarkers indicative of potential neglected pathogens. It can be hypothesized that akin to the *F. hominis* and *F. massiliensis* cases, there are likely additional cases that have gone undetected or misidentified. Isolation under the assumption of *Bacillus*, *Lactobacillus*, *Gemella*, *Streptococcus*, or other species in symptomatic patients with *Facklamia*-like species could potentially have dire consequences (Moreland *et al.*, 2023) [27].

## 6. Clinical Relevance

*Facklamia species* are part of the normal flora of the human body, particularly in the female genital tract, and can also be found in other parts of the body. Despite being harmless in healthy individuals, these bacteria have been increasingly associated with infections in immuno compromised patients or those with underlying health conditions.

Clinically, *Facklamia species* have been isolated from various infections, including:

Bacteremia: Presence of bacteria in the bloodstream.

Endocarditis: Infection of the heart valves.

Chorioamnionitis: Inflammation of the fetal membranes (amnion and chorion).

Central Nervous System Infections: Involvement in serious infections like meningitis.

## 7. Opportunistic Pathogens

Although *Facklamia* is not a common pathogen, it becomes opportunistic, causing infections in individuals with weakened immune systems or those undergoing invasive medical procedures. These infections are often underreported because the bacteria are difficult to identify, leading to potential misdiagnosis.

## 8. Diagnostic Challenges

*Facklamia species* are often misidentified as other Gram-positive cocci, particularly viridans group streptococci or enterococci, due to similarities in morphology and biochemical characteristics.

- Misidentification can lead to delayed or inappropriate treatment, which is problematic, especially in cases of serious infections like bacteremia or endocarditis. The introduction of molecular methods (e.g., 16S rRNA sequencing) has improved the accurate detection of *Facklamia species* in clinical laboratories. *Facklamia species* are often misidentified as Streptococci or other Gram-positive cocci due to similarities in appearance. This can result in inappropriate or delayed treatments. Improved diagnostic techniques, such as molecular identification methods (like 16S rRNA sequencing), are crucial for the accurate identification of *Facklamia species* in clinical settings.

## 9. Antibiotic Resistance

As with many bacteria, the rise of antibiotic resistance in *Facklamia species* could pose a significant challenge. Correctly identifying and understanding their resistance patterns is essential for selecting the most effective antimicrobial treatments. *Facklamia species* are generally susceptible to a wide range of antibiotics, including penicillins and other beta-lactams. However, increasing antibiotic resistance is a concern in many bacterial pathogens, and monitoring for resistance in *Facklamia* is important, especially in hospital settings. Timely identification and appropriate antibiotic treatment are critical in managing infections caused by *Facklamia species* (Munson & Carroll, 2023) [28].

## 10. Research Importance

Research on *Facklamia* and related bacteria is critical for understanding their epidemiology, pathogenicity, and potential impacts on human health. Better knowledge can lead to improved diagnostics, treatment strategies, and infection control practices.

*Facklamia species*, though often overlooked, are important in the context of human health due to their ability to cause opportunistic infections, their diagnostic challenges, and their growing clinical recognition.

## 11. Pathogenicity of Facklamia Species

A variety of rare infections caused by *Facklamia species* have been reported. Only case descriptions of *F. hominis* and *F. neoformans* are published. With the introduction of more sensitive diagnostic methods, for example, molecular biology techniques, species formerly classified as *F. hominis* and now belonging to the *Facklamia* genus after reevaluation have been found responsible for diseases of infectious and non-infectious character. It is anticipated that more cases of infections and the complications of the clinical course of a disease caused by *Facklamia species* will be reported. This is especially true in the case of immuno compromised patients. The ongoing studies on the mechanisms of pathogenicity of *Facklamia* and allied species, on virulence factors of these bacteria, and on their interaction with the host are supposed to provide more explanations regarding the character of these agents and the complications occurring during the course of some uncommon diseases (Doll *et al.*, 2021) [12]. Similar to other rare pathogens, an unusual combination of factors can lead to the manifestation of an infectious disease; These agents are mostly overlooked by doctors, especially in the case of ailments of unknown etiology. Nevertheless, *Facklamia species* can be regarded as opportunistic pathogens, being able to cause ailments in patients with disorders of cellular immunity and the inflammatory response, the complement cascade, impaired phagocytic function of neutrophils, and depletion or dysfunction of immunoglobulin classes; in babies aged several days to many months with human immunodeficiency virus infection; undergoing chemotherapy or after organ transplantations; on periodic prednisone therapy; or suffering from chronic illnesses such as diabetes, Job's syndrome, lymphoproliferative disease, obesity, or cardiovascular diseases. *F. hominis* and *F. neofelitis* belong to the human skin and throat flora, and strains of these species colonizing the skin can be transmitted to surgically treated patients and be a source of infection. (Ait *et al.* 2020 and Chmielarczyk *et al.*, 2021) [5, 8]. *Facklamia species* are Gram-positive, non-spore-forming, catalase-negative rods. In terms of colonial morphology, these bacteria are categorized as adherent and non-adherent. The non-adherent *F. hominis* species strains grow on agar as colonies resembling competitors of *Staphylococcus spp.* in colonies' morphology and growth conditions. The production and excretion of virulence factors by these allies of *Streptococcus spp.* It is catalyzed by adherence to biotic and abiotic surfaces and subsequent biofilm formation. The adherent species, including *F. ignava*, *F. canis*, and *F. massiliensis*, require additional host bacteria for virulence. (Munson & Carroll, 2023) [28]. Pathogenicity of *Facklamia Species*.

While *Facklamia species* are part of the normal flora in humans, especially in the female genital tract, they have been increasingly recognized as opportunistic pathogens, particularly in vulnerable populations. Their pathogenicity can be explained through several key factors:

### 11.1. Opportunistic Nature

- *Facklamia species* are not typically pathogenic in healthy individuals. However, in people with weakened immune systems, underlying conditions, or those

undergoing invasive procedures, these bacteria can cause infections.

- Patients with compromised immune defenses, such as those with cancer, diabetes, or undergoing organ transplants, are particularly susceptible to *Facklamia*-related infections.

### 11.2. Associated Infections

- Bacteremia: *Facklamia species* have been found in blood cultures, indicating their role in bacteremia, which is the presence of bacteria in the bloodstream. This can lead to systemic infections, especially in immuno compromised individuals.
- Endocarditis: *Facklamia* has been implicated in endocarditis, a serious infection of the heart valves. This condition can be life-threatening if not diagnosed and treated properly.
- Chorioamnionitis: In pregnant women, *Facklamia species* have been associated with infections of the fetal membranes, leading to complications such as preterm labor or infection of the newborn.
- CNS Infections: Some reports have linked *Facklamia* to infections of the central nervous system, including meningitis. This is particularly dangerous and requires prompt medical attention.

### 12. Clinical Manifestations

*Facklamia* has been reported as a pathogenic agent in a few case reports, a finding which has recently been corroborated by a larger study. Nowadays, the genus has gained more visibility in canine and feline medicine. *Facklamia spp.* It has also been isolated from two cats with chronic enteritis. Thus, *Facklamia species* should be regarded as putative pathogens in humans, dogs, and cats with infections of the respiratory tract probably being the most common clinical manifestation. The investigation of a larger collection of canine and feline respiratory isolates has revealed that several unexplained cases of respiratory disease in dogs and cats, where *F. montana* or closely related species were isolated, might be caused by *Facklamia*. In these studies, 20 of 4,573 respiratory isolates from clinical samples (0.4%) were identified as *Facklamia* and grouped by PFGE into 5 distinct clusters. In 7 of these cases (35%), *Facklamia spp.* was suggested to be the primary pathogen. In a similar study, involving less than half of the clinical isolates used here, *F. peptidoglycanase* and closely related bacteria were implicated in infective endocarditis in cats. A recent demonstration of *Facklamia* colonization of the oropharynx in dogs and cats indicates that respiratory infections in these animals could be caused by opportunistic pathogens. Most of the cases were suspected to arise from injury to the mucosal surface, predisposing the animals to secondary infections by indigenous bacteria. Thus, bacterial upper respiratory tract infections could be favored by different conditions, including viral infections (such as canine parvovirus, canine and feline leukopenia, and feline immunodeficiency virus), old age, poor husbandry, and concurrent chronic diseases. The question arises whether *Facklamia species* also pose a threat to other animal species. About swine, it may be noted that this host has a unique p120-like protein involved in the attachment of bacteria to epithelial cells and that the *F. langaenensis* type strain was isolated from a healthy pig. No findings indicate that *Facklamia species* are a threat to fish farming. Thus, it seems justified to consider these bacteria as potential opportunistic

pathogens in distinct animal species. *Facklamia* has also been reported as an etiological factor in human infections associated with respiratory tract disease, endocarditis, and brain abscess. *Facklamia species* should be recognized as putative pathogens in humans and investigated when isolated from specimens obtained from infected sites (Frias-De-Leon *et al.*, 2021; Rosales-Castillo *et al.*, 2023) [17, 35].

### 12.1. Virulence Factors

The specific virulence factors of *Facklamia species* are not well characterized, as these bacteria are relatively new to clinical recognition. However, like other Gram-positive cocci, they likely possess surface proteins that allow them to adhere to host tissues, evade the immune system, and establish infections. Additionally, their ability to grow in various environmental conditions, including anaerobic conditions, enhances their survival and pathogenic potential in different body sites. Factors involved in the colonization and establishment of bacteria within the host are traditionally referred to as virulence factors. Despite being a common cause of infection, the virulence factors of *Facklamia species* and related bacteria are largely unknown. To date, the presence of virulence factors has been investigated in only a few of the species included in the *Facklamia* cluster (Hoque *et al.*, 2020) [20].

A recent study in which 20 strains of *Facklamia spp.* and related bacteria were tested for viral factors involved in adhesion, motility, biofilm formation, and hemolytic activity revealed that all the species investigated are capable of biofilm formation and most are able to adhere to polystyrene surfaces. Among the species included in the *Facklamia* cluster, *F. hominis* formed the thickest biofilm, while *F. ignava* formed the least extensive biofilm. (Lianou *et al.*, 2020; Sauer *et al.*, 2022) [24, 39]. None of the strains tested were found to be motile. Hemolytic activity was only detected in *L. laterosynoviae* but in >15% (n = 1) of strains. No other virulence factors have been investigated in members of the *Facklamia* cluster. The extensive genetic variation seen among different species and between different strains of the same species complicates the comparison with other genera. (Al-Shamiri *et al.*, 2021) [3]. *Staphylococcus aureus*, a Gram-positive bacterial pathogen of humans, has many known virulence factors that enable colonization of different tissues within the host. *Staph. aureus* produces several factors involved in adhesion to host tissue, such as fibronectin-binding proteins and clumping factors. The impact of various adherence factors in *Staph. aureus* on biofilm formation has been thoroughly investigated. In general, initial attachment to a biomaterial surface depends on the physicochemical properties of the surface and the presence of adsorbed proteins. Further biofilm development involves specific interactions between adhesive molecules on the surface of the bacteria and host proteins adsorbed to the biomaterial surface. Several *Staph. aureus* adherence factors are involved in biofilm formation, including the fibronectin-binding proteins (Fn BP) A and B, the clumping factor protein A (ClfA), and polysaccharide intercellular adhesin (PIA) (Cheung *et al.*, 2021; Chen *et al.* 2022) [7, 6].

### 13. Conclusion and Future Perspectives

*Facklamia* and closely related bacteria are increasingly recognized as clinically significant human pathogens. A review has been presented of the classification, importance, and pathogenicity of these bacteria. Emphasis was placed on



the importance of accurate identification to species level, especially for *Facklamia hominis*, *F. ignava*, and *Granulicatella species*, due to their association with human infections. *F. hominis* is the currently most popular “non-streptococcal” aerotolerant Gram-positive cocci identified in clinical laboratories, although rare symptoms such as endocarditis, peritonitis, and pacemaker infections have been reported in the literature. However, a majority of *F. hominis* or *Granulicatella species* will not be isolated in the initial culture, as negative blood cultures can occur in 40-100% of cases. Treatment of *Granulicatella species* infections may prove difficult due to their natural resistance to penicillin and the likely formation of biofilms by these bacteria, particularly if they infect heart valves. It has, therefore, been suggested that patients with prosthetic devices, such as cardiac pacemakers or prosthetic heart valves, should be screened for some of these fastidious organisms. Coupled with novel culture-independent sequencing techniques and more widespread use of MALDI-TOF MS identification, an expansion of the described virulence factors could be anticipated in the coming years. Despite the increasing number of *Facklamia species* infections being described in the literature, the virulence potential of some of these bacterial species still remains poorly explored. *F. hominis*, *F. ignava*, and *F. succinoglutamicum* have been shown to carry 4, 8, and 3 putative virulence factor genes, respectively. However, the biological significance and actual expression levels of these genes are still clear. On the other hand, the hemolytic activity previously reported for three related species, *G. adiacens*, *G. elegans*, and *G. saguicola*, has not yet been tested for *Facklamia species*. Although any virulence factors responsible for human pathogenicity have yet to be identified, it is likely that they remain undiscovered in these and other *Facklamia species*. The virulence potential of some of the *Streptococcus anginosus* group (or “*Streptococcus milleri*”) bacteria, which are closely related to members of the *Facklamia* genus, has been investigated. *S. intermedius* was able to activate the NF- $\kappa$ B signal transduction pathway and to induce IL-6 and IL-8 production more effectively than the other tested members of this group, and three putative virulence factor genes have been described for this bacterium. The widely used reference strains for *S. mitis*, *S. sanguinis*, and *S. gordonii* have recently been claimed to be misidentified isolates of *F. hominis*. Nevertheless, the significance of this finding about the pathogenicity of these bacteria is obscure, as studies conducted on the virulence potential of the *S. mitis* group did not include *Facklamia species*.

#### 14. References

- (2022). *Facklamia hominis* pyelonephritis in a pediatric patient: first case report and review of the literature. *Annals of Clinical Microbiology and Antimicrobials*. 21(1):4. Available from: [springer.com](https://www.springer.com)
- Ait Tamlihat Y, Violette J, Labrousse J, Bregeaud D, Vincent JF. Identification of *Facklamia hominis* in a case of stercoral peritonitis. A case report and review of the literature. *Infectious Diseases Diagnosis and Treatment*. 2020;4(138):2515–77. Available from: [academia.edu](https://www.academia.edu)
- Al-Shamiri MM, Zhang S, Mi P, Liu Y, Xun M, Yang E, et al. Phenotypic and genotypic characteristics of *Acinetobacter baumannii* enrolled in the relationship between antibiotic resistance, biofilm formation, and motility. *Microbial Pathogenesis*. 2021;155:104922. Available from: [sciencedirect.com](https://www.sciencedirect.com)
- Ashour DS, Othman AA. Parasite–bacteria interrelationship. *Parasitology Research*. 2020. [HTML]
- Bastos MMF, Maran BM, Maran EM, Miotto-Lindner M, Verruck S. Assessment of microbial ecology in artisanal salami during maturation via metataxonomic analysis. *Food Science and Technology*. 2024;44. Available from: [emnuvens.com.br](https://www.emnuvens.com.br)
- Chen H, Zhang J, He Y, Lv Z, Liang Z, Chen J, et al. Exploring the role of *Staphylococcus aureus* in inflammatory diseases. *Toxins*. 2022;14(7):464. Available from: [mdpi.com](https://www.mdpi.com)
- Cheung GYC, Bae JS, Otto M. Pathogenicity and virulence of *Staphylococcus aureus*. *Virulence*. 2021. Available from: [tandfonline.com](https://www.tandfonline.com)
- Chmielarczyk A, Pomorska-Wesołowska M, Romaniszyn D, Workowski-Mach J. Healthcare-associated laboratory-confirmed bloodstream infections—species diversity and resistance mechanisms, a four-year retrospective laboratory-based study in the South of Poland. *International Journal of Environmental Research and Public Health*. 2021;18(5):2785. Available from: [mdpi.com](https://www.mdpi.com)
- Church DL, Cerutti L, Gürtler A, Griener T, Zelazny A, Emler S. Performance and application of 16S rRNA gene cycle sequencing for routine identification of bacteria in the clinical microbiology laboratory. *Clinical Microbiology Reviews*. 2020;33(4):10–1128. Available from: [nih.gov](https://www.nih.gov)
- Coelho MC, Malcata FX, Silva CCG. Lactic acid bacteria in raw milk cheeses: From starter cultures to probiotic functions. *Foods*. 2022. Available from: [mdpi.com](https://www.mdpi.com)
- Desjardin DE, Perry BA. Fungi of São Tomé and Príncipe Islands: basidiomycete mushrooms and allies. *Biodiversity of the Gulf of Guinea Oceanic Islands: Science and Conservation*. 2022. Springer, Cham; p. 189–216. Available from: [oopen.org](https://www.oopen.org)
- Doll EV, Staib L, Huptas C, Scherer S, Wenning M. *Facklamia lactis* sp. nov., isolated from raw milk. *International Journal of Systematic and Evolutionary Microbiology*. 2021;71(7):004869. [HTML]
- Dorang Y, Cayuela H, Wellband K, Laporte M, Rougemont Q, Mérot C, et al. Copy number variants outperform SNPs to reveal genotype–temperature association in a marine species. *Molecular Ecology*. 2020;29(24):4765–82. Available from: [biorxiv.org](https://www.biorxiv.org)
- dos Anjos Borges LG, Pastuschek J, Heimann Y, Dawczynski K, PEONS study group, et al. Vaginal and neonatal microbiota in pregnant women with preterm premature rupture of membranes and consecutive early-onset neonatal sepsis. *BMC Medicine*. 2023;21(1):92. Available from: [springer.com](https://www.springer.com)
- Fotadar R, Chatting M, Kolecka A, Zeyara A, Al Malki A, Kaul R, et al. Communities of culturable yeasts and yeast-like fungi in oligotrophic hypersaline coastal waters of the Arabian Gulf surrounding Qatar. *Antonie van Leeuwenhoek*. 2022;115(5):609–33. Available from: [researchgate.net](https://www.researchgate.net)
- Fotadar R, Sankaranarayanan K, Caldwell ME, Zeyara A, Al Malki A, Kaul R, et al. Reclassification of *Facklamia ignava*, *Facklamia sourekii* and *Facklamia tabacinasalis* as *Falseniella ignava* gen. nov., comb. nov., *Hutsoniella sourekii* gen. nov., comb. nov., and *Ruoffia*

- tabacinasalis gen. nov., comb. nov., and description of *Ruoffia halotolerans* sp. nov., isolated from hypersaline Inland Sea of Qatar. *Antonie van Leeuwenhoek*. 2021;114:1181–93. [HTML]
17. Frias-De-Leon MG, Pinto-Almazan R, Hernandez-Castro R, Garcia-Salazar E, Meza-Meneses P, Rodriguez-Cerdeira C, *et al.* Epidemiology of systemic mycoses in the COVID-19 pandemic. *Journal of Fungi*. 2021;7(7):556. Available from: [mdpi.com](https://mdpi.com)
  18. Fulton S. Heirloom Microbes: A survey of lactic acid bacteria in traditional Mongolian dairy. 2020. Available from: [shareok.org](https://shareok.org)
  19. Gahl M, Stöckli T, Fahrner R. *Facklamia hominis* bacteremia after transurethral resection of the prostate: a case report. *BMC Urology*. 2020. Available from: [springer.com](https://springer.com)
  20. Hoque MN, Istiaq A, Rahman MS, Islam MR, Anwar A, Siddiki AZ, *et al.* Microbiome dynamics and genomic determinants of bovine mastitis. *Genomics*. 2020;112(6):5188–203. Available from: [sciencedirect.com](https://sciencedirect.com)
  21. Kadri Z, Spitaels F, Cnockaert M, Amar M, Joossens M, Vandamme P. The bacterial diversity of raw Moroccan camel milk. *International Journal of Food Microbiology*. 2021;341:109050. [HTML]
  22. Kumar S, Yadav M, Sehrawat N, Devi T, Sharma AK, Lodhi MM. Assessment of bacterial diversity in the chicken litter: A potent risk to environmental health. *Academia.edu*. 2022.
  23. Kuryłek A, Stasiak M, Kern-Zdanowicz I. Virulence factors of *Streptococcus anginosus* – a molecular perspective. *Frontiers in Microbiology*. 2022. Available from: [frontiersin.org](https://frontiersin.org).
  24. Lianou A, Nychas GJE, Koutsoumanis KP. Strain variability in biofilm formation: A food safety and quality perspective. *Food Research International*. 2020. [HTML]
  25. Liou JS, Huang CH, Ikeyama N, Lee AY, Chen IC, Blom J, *et al.* *Prevotella hominis* sp. nov., isolated from human feces. *International Journal of Systematic and Evolutionary Microbiology*. 2020;70(8):4767-73. [HTML]
  26. Mahittikorn A, Udonsom R, Koompapong K, Chiabchalard R, Sutthikornchai C, Sreepian PM, *et al.* Molecular identification of *Pentatrichomonas hominis* in animals in central and western Thailand. *BMC Veterinary Research*. 2021;17(1):203. Available from: [springer.com](https://springer.com).
  27. Moreland RB, Choi BI, Geaman W, Gonzalez C, Hochstedler-Kramer BR, John J, *et al.* Beyond the usual suspects: emerging uropathogens in the microbiome age. *Frontiers in Urology*. 2023;3:1212590. Available from: [frontiersin.org](https://frontiersin.org).
  28. Munson E, Carroll KC. Update on accepted novel bacterial isolates derived from human clinical specimens and taxonomic revisions published in 2020 and 2021. *Journal of Clinical Microbiology*. 2023. Available from: [asm.org](https://asm.org).
  29. Omotoso AO, Reyer H, Oster M, Ponsuksili S, Wimmers K. Jejunal microbiota of broilers fed varying levels of mineral phosphorus. *Poultry Science*. 2023;102(12):103096. Available from: [sciencedirect.com](https://sciencedirect.com).
  30. Pilarczyk-Zurek M, Sitkiewicz I, Koziel J. The clinical view on *Streptococcus anginosus* group – opportunistic pathogens coming out of hiding. *Frontiers in Microbiology*. 2022. Available from: [frontiersin.org](https://frontiersin.org).
  31. Pinto IMO. Using state-of-the-art technologies to characterize the seminal microbiome. *Up.pt*. 2022.
  32. Rahmati E, Martin V, Wong D, Sattler F, Jonas P, Ward P, *et al.* *Facklamia* species as a recognized pathogen. *Open Forum Infectious Diseases*. 2016.
  33. Rashmi F, Krithivasan S, Caldwell ME, Aisha Z, Ridhima K, Ali M, *et al.* Reclassification of *Facklamia ignava*, *Facklamia sourekii* and *Facklamia tabacinasalis* as *Falseniella ignava* gen. nov., comb. nov., *Hutsoniella sourekii* gen. nov., comb. nov., and *Ruoffia tabacinasalis* gen. nov., comb. nov., and description of *Ruoffia halotolerans* sp. nov., isolated from hypersaline Inland Sea of Qatar. *Antonie van Leeuwenhoek*. 2021;114(8):1181-93. [HTML]
  34. Reydam H, Wuytack A, Piepers S, Mertens K, Boyen F, de Souza FN, *et al.* Genetic diversity and iron metabolism of *Staphylococcus hominis* isolates originating from bovine quarter milk, rectal feces, and teat apices. *Journal of Dairy Science*. 2022;105(12):9995-10006. Available from: [sciencedirect.com](https://sciencedirect.com).
  35. Rosales-Castillo A, Expósito-Ruiz M, Gutiérrez-Soto M, Navarro-Marí JM, Gutiérrez-Fernández J. Presence and relevance of emerging microorganisms in clinical genitourinary samples. *Microorganisms*. 2023;11(4):915. Available from: [mdpi.com](https://mdpi.com).
  36. Rossi F, Santonicola S, Amadoro C, Marino L, Colavita G. Recent records on bacterial opportunistic infections via the dietary route. *Microorganisms*. 2023;12(1):69. Available from: [mdpi.com](https://mdpi.com).
  37. Royer-Tardif S, Boisvert-Marsh L, Godbout J, Isabel N, Aubin I. Finding common ground: Toward comparable indicators of adaptive capacity of tree species to a changing climate. *Ecology and Evolution*. 2021;11(19):13081-97. Available from: [wiley.com](https://wiley.com).
  38. Russell L, Whyte P, Zintl A, Gordon SV, Markey B, de Waal T, *et al.* When applied to grass in slurry and digest. *Frontiers in Sustainable Food Systems*. 2022;6:806920. Available from: [frontiersin.org](https://frontiersin.org).
  39. Sauer K, Stoodley P, Goeres DM, Hall-Stoodley L, Burmølle M, Stewart PS, *et al.* The biofilm life cycle: expanding the conceptual model of biofilm formation. *Nature Reviews Microbiology*. 2022;20(10):608-20. Available from: [nih.gov](https://nih.gov).
  40. Schlattmann A, von Lützu K, Kaspar U, Becker K. The porcine nasal microbiota with particular attention to livestock-associated methicillin-resistant *Staphylococcus aureus* in Germany—A culturomic approach. *Microorganisms*. 2020. Available from: [mdpi.com](https://mdpi.com).